

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00	A2	(11) International Publication Number: WO 98/51787 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/GB98/01294 (22) International Filing Date: 5 May 1998 (05.05.98) (30) Priority Data: 9709421.3 10 May 1997 (10.05.97) GB (71) Applicant (for all designated States except US): ZENECA LIMITED (GB/GB); 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): EMERY, Stephen, Charles [GB/GB]; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). BLAKEY, David, Charles [GB/GB]; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (74) Agent: GILES, Allen, Frank; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: CHEMICAL COMPOUNDS (57) Abstract The invention provides a gene construct encoding a cell targeting moiety and a heterologous prodrug activating enzyme for use as a medicament in a mammalian host wherein the gene construct is capable of expressing the cell targeting moiety and enzyme as a conjugate within a target cell in the mammalian host and wherein the conjugate is directed to leave the cell thereafter for selective localisation at a cell surface antigen recognised by the cell targeting moiety.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Licchtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CHEMICAL COMPOUNDS

This invention relates particularly to gene directed enzyme prodrug therapy (GDEPT) using *in situ* antibody generation to provide enhanced selectivity, particularly for use in cancer
5 therapy.

Known gene therapy based prodrug therapeutic approaches include virus-directed enzyme prodrug therapy (VDEPT) and gene-directed enzyme prodrug therapy (GDEPT), the latter term encompassing both VDEPT and non-viral delivery systems. VDEPT involves targeting tumour cells with a viral vector carrying a gene which codes for an enzyme capable
10 of activating a prodrug. The viral vector enters the tumour cell and enzyme is expressed from the enzyme gene inside the cell. In GDEPT, alternative approaches such as microinjection, liposomal delivery and receptor mediated DNA uptake as well as viruses may be used to deliver the gene encoding the enzyme.

In both VDEPT and GDEPT the enzyme gene can be transcriptionally regulated by
15 DNA sequences capable of being selectively activated in mammalian cells e.g. tumour cells (EP 415 731 (Wellcome); Huber *et al*, Proc. Natl. Acad. Sci. USA, 88, 8039-8043, 1991). While giving some degree of selectivity, gene expression may also occur in non-target cells and this is clearly undesirable when the approach is being used to activate prodrugs into potent cytotoxic agents. In addition these regulatory sequences will generally lead to reduced
20 expression of the enzyme compared with using viral promoters and this will lead to a reduced ability to convert prodrug in the target tissue.

Expression and localisation of the prodrug activating enzyme inside the cell has disadvantages. Prodrug design is severely limited by the fact that the prodrug has to be able to cross the cell membrane and enter the cell but not be toxic until it is converted to the drug
25 inside the cell by the activating enzyme. Most prodrugs utilise hydrophilic groups to prevent cell entry and thus reduce cytotoxicity. Prodrug turnover by activating enzyme produces a less hydrophilic drug which can enter cells to produce anti-cancer effects. This approach can not be used when the activating enzyme is expressed inside the cell. Another disadvantage is that target cells which lack intracellular activating enzyme will be difficult to attack because
30 they are unable to generate active drug. To achieve this desirable "bystander activity" (or "neighbouring cell kill"), the active drug will have to be capable of diffusion out of the cell containing activating enzyme to reach target cells which lack enzyme expression. Many

active drugs when produced inside a cell will be unable to escape from the cell to achieve this bystander effect.

Modifications of GDEPT have been put forward to overcome some of the problems described above. Firstly vectors have been described which are said to express the activating enzyme on the surface of the target cell (WO 96/03515) by attaching a signal peptide and transmembrane domain to the activating enzyme. The approach, if viable, would overcome the problems of having the activating enzyme located inside the cell but would still have to rely on transcriptionally regulated sequences capable of being selectively expressed in target cells to restrict cell expression. As described above there are disadvantages of using such sequences. Secondly vectors have been described which result in secretion of the enzyme from the target cell (WO 96/16179). In this approach the enzyme would be able to diffuse away from its site of generation since it is extracellular and not attached to the cell surface. Enzyme which has diffused away from the target site would be capable of activating prodrug at non-target sites leading to unwanted toxicity. To achieve some selectivity it is suggested that enzyme precursors could be used which are cleaved by pathology associated proteases to form active enzyme. Some selectivity is likely to be achieved by this approach but its unlikely that activation will only occur at target sites. In addition, once activated, the enzyme will still be free to diffuse away from the target site and thus suffer from the same drawback described above.

For GDEPT approaches, three levels of selectivity can be observed. Firstly, there is selectivity at the cell infection stage such that only specific cell types are targeted. For example cell selectivity can be provided by the gene delivery system *per se*. An example of this type of selectivity is set out in International Patent Application WO 95/26412 (UAB Research Foundation) which describes the use of modified adenovirus fiber proteins incorporating cell specific ligands. Other examples of cell specific targeting include *ex vivo* gene transfer to specific cell populations such as lymphocytes and direct injection of DNA into muscle tissue.

The second level of selectivity is control of gene expression after cell infection such as for example by the use of cell or tissue specific promoters. If the gene has been delivered to a cell type in a selective manner then it is important that a promoter is chosen that is compatible with activity in the cell type.

- 3 -

The third level of selectivity can be considered as the selectivity of the expressed gene construct. Selectivity at this level has received scant attention to date. In International patent application WO 96/16179 (Wellcome Foundation) it is suggested that enzyme precursors could be used which are cleaved by pathology associated proteases to form active enzyme.

5 Some selectivity is likely to be achieved by this approach but it is unlikely that activation will only occur at target sites. In addition, once activated, the enzyme will still be free to diffuse away from the target site and thus suffer from the same drawback of activating prodrug at non-target sites leading to unwanted toxicity.

There exists a need for more selective GDEPT systems to reduce undesirable effects in

10 normal tissues arising from erroneous prodrug activation.

The present invention is based on the discovery that antibody-heterologous enzyme gene constructs can be expressed intracellularly and used in GDEPT systems (or other systems such as AMIRACS - see below) for cell targeting arising from antibody specificity to deliver cell surface available enzyme in a selective manner. This approach may be used

15 optionally in combination with any other suitable specificity enhancing technique(s) such as targeted cell infection and/or tissue specific expression.

According to one aspect of the present invention there is provided a gene construct encoding a cell targeting antibody and a heterologous enzyme for use as a medicament in a mammalian host wherein the gene construct is capable of expressing the antibody and enzyme

20 as a conjugate within a target cell in the mammalian host and wherein the conjugate can leave the cell thereafter for selective localisation at a cell surface antigen recognised by the antibody.

According to another aspect of the present invention there is provided a gene construct encoding a cell targeting moiety and a heterologous prodrug activating enzyme for use as a

25 medicament in a mammalian host wherein the gene construct is capable of expressing the cell targeting moiety and heterologous prodrug activating enzyme as a conjugate within a cell in the mammalian host and wherein the conjugate is directed to leave the cell thereafter for selective localisation at a cell surface antigen recognised by the cell targeting moiety.

The "cell targeting moiety" is defined as any polypeptide or fragment thereof which

30 selectively binds to a particular cell type in a host through recognition of a cell surface antigen. Preferably the cell targeting moiety is an antibody. Cell targeting moieties other than antibodies include ligands as described for use in Ligand Directed Enzyme Prodrug

- 4 -

Therapy as described in International patent application WO 97/26918, Cancer Research Campaign Technology Limited, such as for example epidermal growth factor, heregulin, c-erbB2 and vascular endothelial growth factor with the latter being preferred.

A "cell targeting antibody" is defined as an antibody or fragment thereof which
5 selectively binds to a particular cell type in a host through recognition of a cell surface antigen. Preferred cell targeting antibodies are specific for solid tumours, more preferably colorectal tumours, more preferably an anti-CEA antibody, more preferably antibody A5B7 or 806.077 antibody with 806.077 antibody being especially preferred. Hybridoma 806.077 antibody was deposited at the European Collection of Animal Cell Cultures (ECACC), PHLS
10 Centre for Applied Microbiology & Research, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom on 29th February 1996 under accession no. 96022936 in accordance with the Budapest Treaty.

Antibody A5B7 binds to human carcinoembryonic antigen (CEA) and is particularly suitable for targeting colorectal carcinoma. A5B7 is available from DAKO Ltd., 16 Manor
15 Courtyard, Hughenden Avenue, High Wycombe, Bucks HP13 5RE, England, United Kingdom. In general the antibody (or antibody fragment) - enzyme conjugate should be at least divalent, that is to say capable of binding to at least 2 tumour associated antigens (which may be the same or different). Antibody molecules may be humanised by known methods such as for example by "CDR grafting" as disclosed in EP239400 or by grafting complete
20 variable regions from for example a murine antibody onto human constant regions ("chimaeric antibodies") as disclosed in US 4816567. Humanised antibodies may be useful for reducing immunogenicity of an antibody (or antibody fragment). A humanised version of antibody A5B7 has been disclosed in International Patent Application WO 92/01059 (Celltech).

25 The hybridoma which produces monoclonal antibody A5B7 was deposited with the European Collection of Animal Cell Cultures, Division of Biologics, PHLS Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom. The date of deposit was 14th July 1993 and the accession number is No. 93071411. Antibody A5B7 may be obtained from the deposited hybridoma using standard
30 techniques known in the art such as documented in Fenge C, Fraune E & Schuegerl K in "Production of Biologicals from Animal Cells in Culture" (Spier RE, Griffiths JR & Meignier B, eds) Butterworth-Heinemann, 1991, 262-265 and Anderson BL & Gruenberg ML

- 5 -

in "Commercial Production of Monoclonal Antibodies" (Seaver S, ed), Marcel Dekker, 1987, 175-195. The cells may require re-cloning from time to time by limiting dilution in order to maintain good levels of antibody production.

A "heterologous enzyme" is defined as an enzyme for turning over a substrate that has
5 been administered to the host and the enzyme is not naturally present in the relevant compartment of the host. The enzyme may be foreign to the mammalian host (e.g. a bacterial enzyme like CPG2) or it may not naturally occur within the relevant host compartment (e.g. the use of lysozyme as an ADEPT enzyme (for an explanation of ADEPT see below) is possible because lysozyme does not occur naturally in the circulation, see US 5433955, Akzo
10 NV). The relevant host compartment is that part of the mammalian host in which the substrate is distributed. Preferred enzymes are enzymes suitable for ADEPT or AMIRACS (Antimetabolite with Inactivation of Rescue Agents at Cancer Sites; see Bagshawe (1994) in Cell Biophysics 24/25, 83-91) but ADEPT enzymes are preferred. Antibody directed enzyme prodrug therapy (ADEPT) is a known cancer therapeutic approach. ADEPT uses a tumour
15 selective antibody conjugated to an enzyme. The conjugate is administered to the patient (usually intravenously), allowed to localise at the tumour site(s) and clear from the blood and other normal tissues. A prodrug is then administered to the patient which is converted by the enzyme (localised at the tumour site) into a cytotoxic drug which kills the tumour cells.

In International Patent Application WO 96/20011, published 4-Jul-96, we proposed a
20 "reversed polarity" ADEPT system based on mutant human enzymes having the advantage of low immunogenicity compared with for example bacterial enzymes. A particular host enzyme was human pancreatic CPB (see for example, Example 15 [D253K]human CPB & 16 [D253R]human CPB therein) and prodrugs therefor (see Examples 18 & 19 therein). The host enzyme is mutated to give a change in mode of interaction between enzyme and prodrug in
25 terms of recognition of substrate compared with the native host enzyme. In our subsequent International Patent Application No PCT/GB96/01975 (published 6-Mar-97 as WO 97/07796) further work on mutant CPB enzyme/ prodrug combinations for ADEPT are described. Preferred enzymes suitable for ADEPT are any one of CPG2 or a reversed polarity CPB enzyme, for example any one of [D253K]HCPB, [G251T,D253K]HCPB or
30 [A248S,G251T,D253K]HCPB. A preferred form of CPG2 is one in which the polypeptide glycosylation sites have been mutated so as to prevent or reduce glycosylation on expression in mammalian cells (see WO 96/03515, Cancer Research Campaign Technology); this gives

improved enzyme activity. Further considerations arise for enzymes such as CPB which require a pro domain to facilitate correct folding; here the pro domain can either be expressed as a separately (in trans) or expressed as part of the fusion protein and subsequently removed.

Large scale purification of CPG2 from *Pseudomonas* RS-16 was described in

- 5 Sherwood *et al* (1985), Eur. J. Biochem., 148, 447 - 453. CPG2 may be obtained from Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom. CPG2 may also be obtained by recombinant techniques. The nucleotide coding sequence for CPG2 has been published by Minton, N.P. *et al.*, Gene, 31 (1984), 31-38.
- Expression of the coding sequence has been reported in *E.coli* (Chambers, S.P. *et al.*, Appl. Microbiol, Biotechnol. (1988), 29, 572-578) and in *Saccharomyces cerevisiae* (Clarke, L. E. *et al.*, J. Gen Microbiol, (1985) 131, 897-904). Total gene synthesis has been described by M. Edwards in Am. Biotech. Lab (1987), 5, 38-44. Expression of heterologous proteins in *E.coli* has been reviewed by F.A.O. Marston in DNA Cloning Vol. III, Practical Approach Series, IRL Press (Editor D M Glover), 1987, 59-88. Expression of proteins in yeast has
- 15 been reviewed in Methods in Enzymology Volume 194, Academic Press 1991, Edited by C. Guthrie and G R Fink.

Whilst cancer therapeutic approaches are preferred the invention may also be applied to other therapeutic areas as long as a target antigen can be selected and a suitable enzyme/prodrug combination prepared. For example, inflammatory diseases such as rheumatoid

20 arthritis may be treated by for example using an antibody selective for synovial cells fused to an enzyme capable of converting an anti-inflammatory drug in the form of a prodrug into an anti-inflammatory drug. Use of antibodies to target rheumatoid arthritis disease has been described in Blakey *et al*, 1988, Scand. J. Rheumatology, Suppl. 76, 279-287.

A "conjugate" between antibody and enzyme can be a fusion protein (covalent

25 linkage) or the conjugate can be formed by non-covalent binding between antibody and enzyme formed *in situ*. Preferably the conjugate is in the form of a fusion protein, more preferably the antibody component of the fusion is at least divalent (for improved binding avidity compared with monovalent antibody). Antibody constructs lacking an Fc portion are preferred, especially Fab or F(ab')₂ fragments. For CPG2 fusions (or fusions with any non-

30 monomeric enzyme) special considerations apply because CPG2 is a dimeric enzyme and the antibody is preferably divalent thus there exists the potential for undesirable competing dimerisation between two molecular species. Therefore a preferred CPG2 fusion is one in

- 7 -

which the fusion protein is formed through linking a C-terminus of an antibody Fab heavy chain (ie lacking a hinge region) to an N-terminus of a CPG2 molecule; two of these Fab-CPG2 molecules then dimerise through the CPG2 dimerisation domain to form a (Fab-CPG2)₂ conjugate. For antibody constructs with monomeric enzymes, F(ab')₂ fragments are preferred, especially F(ab')₂ fragments having a human IgG3 hinge region. Fusions between antibody and enzyme may optionally be effected through a short peptide linker such as for example (G₄S)₃. Preferred fusion constructs are those in which the enzyme is fused to the C terminus of the antibody, through the heavy or light chain thereof with fusion through the antibody heavy chain being preferred. Accordingly a preferred gene construct is a gene construct for use as a medicament as described herein in which the antibody-enzyme CPG2 conjugate is a fusion protein in which the enzyme is fused to the C terminus of the antibody through the heavy or light chain thereof whereby dimerisation of the encoded conjugate when expressed can take place through a dimerisation domain on CPG2. A more preferred gene construct is a gene construct for use as a medicament wherein the fusion protein is formed through linking a C-terminus of an antibody Fab heavy chain to an N-terminus of a CPG2 molecule to form a Fab-CPG2 whereby two Fab-CPG2 molecules when expressed dimerise through CPG2 to form a (Fab-CPG2)₂ conjugate. In another embodiment of the invention a preferred gene construct for use as a medicament is one wherein the carboxypeptidase is selected from [D253K]HCPB, [G251T,D253K]HCPB or [A248S,G251T,D253K]HCPB.

It is contemplated that should it be possible to obtain a natural multimeric enzyme in monomeric form whilst substantially retaining enzymic activity then the monomeric form of the enzyme could be used to form a conjugate of the invention. Similarly, it is contemplated that should it be possible to obtain a natural monomeric enzyme in multimeric form whilst substantially retaining enzymic activity then the multimeric form of the enzyme could be used to form a conjugate of the invention.

The conjugate is directed to leave the cell after expression therein through use of a secretory leader sequence which is cleaved as the conjugate passes through the cell membrane. Preferably the secretory leader is the secretory leader that occurs naturally with the antibody.

According to another aspect of the present invention there is provided use of a gene construct encoding a cell targeting antibody and a heterologous enzyme for use for manufacture of a medicament for cancer therapy in a mammalian host wherein the gene

construct is capable of expressing the antibody and enzyme as a conjugate within a target cell in the mammalian host and wherein the conjugate can leave the cell thereafter for selective localisation at a cell surface antigen recognised by the antibody.

Any suitable delivery system may be applied to deliver the gene construct of the present invention including viral and non-viral systems. Viral systems include retroviral vectors, adenoviral vectors, adeno-associated virus, vaccinia, herpes simplex virus, HIV, the minute virus of mice, hepatitis B virus and influenza virus. Non-viral systems include uncomplexed DNA, DNA-liposome complexes, DNA-protein complexes and DNA-coated gold particles.

10 Retroviral vectors lack immunogenic proteins and there is no preexisting host immunity but are limited to infecting dividing cells. Retroviruses have been used in clinical trials (Rosenberg *et al.*, N. Engl. J. Med., 1990, 323: 570-578). Retroviruses are composed of an RNA genome that is packaged in an envelope derived from host cell membrane and viral proteins. For gene expression, it must first reverse transcribe its positive-strand RNA genome
15 into double-stranded DNA, which is then integrated into the host cell DNA using reverse transcriptase and integrase protein contained in the retrovirus particle. The integrated provirus is able to use host cell machinery for gene expression.

Murine leukemia virus is widely used (Miller *et al.*, Methods Enzymol., 1993, 217: 581-599). Retroviral vectors are constructed by removal of the gag, pol and env genes to
20 make room for the relevant payload and to eliminate the replicative functions of the virus. Virally encoded mRNAs are eliminated and this removes any potential immune response to the transduced cells. Genes encoding antibiotic resistance often are included as a means of selection. Promoter and enhancer functions also may be included for example to provide for tissue-specific expression after administration *in vivo*. Promoter and enhancer functions
25 contained in the long terminal repeat may also be used.

These viruses can be produced only in viral packaging cell lines. The packaging cell line may be constructed by stably inserting the deleted viral genes (*gag*, *pol*, and *env*) into the cell such that they reside on different chromosomes to prevent recombination. The packaging cell line is used to construct a producer cell line that will generate replication-defective
30 retrovirus containing the relevant payload gene by inserting the recombinant proviral DNA. Plasmid DNA containing the long terminal repeat sequences flanking a small portion of the *gag* gene that contains the encapsidation sequence and the genes of interest is transfected into

the packaging cell line using standard techniques for DNA transfer and uptake (electroporation, calcium precipitation, *etc.*). Variants of this approach have been employed to decrease the likelihood of production of replication-competent virus (Jolly, D., Cancer Gene Therapy, 1994, 1, 51-64). The host cell range of the virus is determined by the
5 envelope gene (*env*) and substitution of *env* genes with different cell specificities can be employed. Incorporation of appropriate ligands into the envelope protein may also be used for targeting.

Administration may be achieved by any suitable technique e.g. *ex vivo* transduction of patients' cells, by the direct injection of virus into tissue, and by the administration of the
10 retroviral producer cells.

The *ex vivo* approach has a disadvantage in that it requires the isolation and maintenance in tissue culture of the patient's cells, but it has the advantage that the extent of gene transfer can be quantified readily and a specific population of cells can be targeted. In addition, a high ratio of viral particles to target cells can be achieved and thus improve the
15 transduction efficiency (Anderson *et al.*, Hum. Gene Ther., 1990, 1: 331-341; Rosenberg *et al.*, N. Engl. J. Med., 1990, 323: 570-578; Culver *et al.*, Hum. Gene Ther., 1991, 2: 107-109; Nienhuis *et al.*, Cancer, 1991, 67: 2700-2704, Anderson *et al.*, Hum. Gene Ther., 1990, 1: 331-341, Grossman *et al.*, Nat. Genet., 1994, 6: 335-341, Lotze *et al.*, Hum. Gene Ther., 1992, 3: 167-177; Lotze, M.T., Cell Transplant., 1993, 2: 33-47; Lotze *et al.*, Hum. Gene
20 Ther., 1994, 5: 41-55 and US patent 5399346 (Anderson). In some cases direct introduction of virus *in vivo* is necessary. Retroviruses have been used to treat brain tumours wherein the ability of a retrovirus to infect only dividing cells (tumour cells) may be particularly advantageous.

To increase efficiency Oldfield *et al.*, in Hum. Gene Ther., 1993, 4: 39-69 proposed
25 the administration of a retrovirus producer cell line directly into patients' brain tumours. The murine producer cell would survive within the brain tumour for a period of days, and would secrete retrovirus capable of transducing the surrounding brain tumour. Virus carrying the herpes virus thymidine kinase gene renders cells susceptible to killing by ganciclovir, which is metabolized to a cytotoxic compound by thymidine kinase. Patent references on retroviruses
30 are: EP 334301, WO 91/02805 & WO 92/05266 (Viagene) and; US 4650764 (University of Wisconsin).

Human adenoviral infections have been described (*see* Horwitz, M.S., In Virology, 2nd ed. Raven Press, New York, 1990, pp. 1723-1740). Most adults have prior exposure to adenovirus and have antiadenovirus antibodies. These viruses possess a double-stranded DNA genome, and replicate independent of host cell division.

5 Adenoviral vectors possess advantageous properties. They are capable of transducing a broad spectrum of human tissues and high levels of gene expression can be obtained in dividing and nondividing cells. Several routes of administration can be used including intravenous, intrabiliary, intraperitoneal, intravesicular, intracranial and intrathecal injection, and direct injection of the target organ. Thus targeting based on anatomical boundaries is
10 feasible.

The adenoviral genome encodes about 15 proteins and infection involves a fiber protein to bind a cell surface receptor. The penton base of the capsid engages integrin receptor domains ($\alpha_3\beta_3$, or $\alpha_3\beta_5$) on the cell surface resulting in internalization of the virus. Viral DNA enters the nucleus and begins transcription without cell division. Expression and
15 replication is under control by the E1A and E1B genes (*see* Horwitz, M.S., In Virology, 2nd ed., 1990, pp. 1723-1740). Removal of E1 genes renders the virus replication-incompetent. Expression of adenoviral proteins leads to both an immune response which may limit effectiveness particularly on repeat administration. However, recent approaches in which other adenoviral genes such as the E2a gene (which controls expression of the fibre knob and
20 a number of other viral proteins) are also removed from the viral genome may abolish or greatly reduce the expression of many of these viral proteins in target cells.

Adenoviral serotypes 2 and 5 have been extensively used for vector construction. Bett *et al.*, Proc. Nat. Acad. Sci. U.S.A., 1994, 91: 8802-8806 have used an adenoviral type 5 vector system with deletions of the E1 and E3 adenoviral genes. The 293 human embryonic
25 kidney cell line has been engineered to express E1 proteins and can thus transcomplement the E1-deficient viral genome. The virus can be isolated from 293 cell media and purified by limited dilution plaque assays (Graham, F.L. and Prevek, L. In Methods in Molecular Biology: Gene Transfer and Expression Protocols, Humana Press 1991, pp. 109-128). Recombinant virus can be grown in 293 cell line cultures and isolated by lysing infected cells
30 and purification by caesium chloride density centrifugation. One problem of the 293 cells for manufacture of recombinant adenovirus is that due to additional flanking regions of the E1 genes is that they may give rise to replication competent adenovirus (RCA) during the viral

- 11 -

particle production. Although this material is only wild type adenovirus and not replication competent recombinant virus it can have significant effects on the eventual yield of the desired adenoviral material and lead to increased manufacturing costs, quality control issues for the production runs and acceptance of batches for clinical use. Alternative cell lines such as the PER.C6 which have more defined E1 gene integration than 293 cells (i.e. contain not flanking viral sequence) have been developed which do not allow the recombination events which produce RCA and thus have the potential to overcome above viral production issues.

Adenoviral vectors have the disadvantage of relatively short duration of transgene expression due to immune system clearance and dilutional loss during target cell division but improvements in vector design are anticipated. Patent references on adenoviruses are: WO 96/03517 (Boehringer); WO 96/13596 (Rhone Poulenc Rorer); WO 95/29993 (University of Michigan) and; WO 96/34969 (Canji). Recent advances in adenoviral vectors for cancer gene therapy including the development of strategies to reduce immunogenicity, chimeric adenoviral/retroviral vectors and conditional (or restricted) replicative recombinant adenoviral systems are reviewed in Bilbao *et al.*, Exp. Opin. Ther. Patents, 1997, 7 (12):1427-1446.

Adeno-associated virus (AAV) (Kotin, R.M., Hum. Gene Ther., 1994, 5: 793-801) are single-stranded DNA, nonautonomous parvoviruses able to integrate into the genome of nondividing cells of a very broad host range. AAV has not been shown to be associated with human disease and does not elicit an immune response.

AAV has two distinct life cycle phases. Wild-type virus will infect a host cell, integrate and remain latent. In the presence of adenovirus, the lytic phase of the virus is induced, which is dependent on the expression of early adenoviral genes, and leads to active virus replication. The AAV genome is composed of two open reading frames (called *rep* and *cap*) flanked by inverted terminal repeat (ITR) sequences. The *rep* region encodes four proteins which mediate AAV replication, viral DNA transcription, and endonuclease functions used in host genome integration. The *rep* genes are the only AAV sequences required for viral replication. The *cap* sequence encodes structural proteins that form the viral capsid. The ITRs contain the viral origins of replication, provide encapsidation signals, and participate in viral DNA integration. Recombinant, replication-defective viruses that have been developed for gene therapy lack *rep* and *cap* sequences. Replication-defective AAV can be produced by cotransfecting the separated elements necessary for AAV replication into a

permissive 293 cell line. Patent references on AAV include: WO 94/13788 (University of Pittsburgh) and US 4797368 (US Department of Health).

Gene therapy vectors from pox viruses have been described (Moss, B. and Flexner, C., *Annu. Rev. Immunol.*, 1987, 5: 305-324; Moss, B., In *Virology*, 1990, pp. 2079-2111).

- 5 Vaccinia are large, enveloped DNA viruses that replicate in the cytoplasm of infected cells. Nondividing and dividing cells from many different tissues are infected, and gene expression from a nonintegrated genome is observed. Recombinant virus can be produced by inserting the transgene into a vaccinia-derived plasmid and transfecting this DNA into vaccinia-infected cells where homologous recombination leads to the virus production. A significant
- 10 disadvantage is that it elicits a host immune response to the 150 to 200 virally encoded proteins making repeated administration problematic.

- The herpes simplex virus is a large, double-stranded DNA virus that replicates in the nucleus of infected cells suitable for gene delivery (*see* Kennedy, P.G.E. and Steiner, I., *Q.J. Med.*, 1993, 86: 697-702). Advantages include a broad host cell range, infection of dividing
- 15 and nondividing cells, and large sequences of foreign DNA can be inserted into the viral genome by homologous recombination. Disadvantages are the difficulty in rendering viral preparations free of replication-competent virus and a potent immune response. Deletion of the viral thymidine kinase gene renders the virus replication-defective in cells with low levels of thymidine kinase. Cells undergoing active cell division (*e.g.*, tumour cells) possess
- 20 sufficient thymidine kinase activity to allow replication. Cantab Pharmaceuticals have a published patent application on herpes viruses (WO 92/05263).

A variety of other viruses, including HIV, the minute virus of mice, hepatitis B virus, and influenza virus, have been considered as possible vectors for gene transfer (*see* Jolly, D., *Cancer Gene Therapy*, 1994, 1: 51-64).

- 25 The use of attenuated *Salmonella Typhimurium* bacteria which specifically target and replicate in hypoxic environments (such as are found in the necrotic centres of tumours) as gene delivery vehicles for prodrug enzyme based therapy (Tumour Amplified Prodrug Enzyme Therapy known as TAPET™) has also been proposed and is under development by Vion Pharmaceuticals. This system offers a further gene delivery alternative to the viral and
- 30 non-viral delivery approaches discussed below.

Nonviral DNA delivery strategies are also applicable. These DNA delivery systems include uncomplexed plasmid DNA, DNA-liposome complexes, DNA-protein complexes,

and DNA-coated gold particles.

Purified nucleic acid can be injected directly into tissues and results in transient gene expression for example in muscle tissue, particularly effective in regenerating muscle (Wolff *et al.*, Science, 1990, 247: 1465-1468). Davis *et al.*, in Hum. Gene Ther., 1993, 4: 733-740
5 has published on direct injection of DNA into mature muscle. Skeletal and cardiac muscle is generally preferred. Patent references are: WO 90/11092, US 5589466 (Vical) and WO 97/05185 (biodegradable DNA impregnated hydrogels for injection, Focal).

Plasmid DNA on gold particles can be "fired" into cells (e.g. epidermis or melanoma) using a gene-gun. DNA is coprecipitated onto the gold particle and then fired using an
10 electric spark or pressurized gas as propellant (Fynan *et al.*, Proc. Natl. Acad. Sci. U.S.A., 1993, 90: 11478-11482). Electroporation has also been used to enable transfer of DNA into solid tumours using electroporation probes employing multi-needle arrays and pulsed, rotating electric fields (Nishi *et al.*, in Cancer Res., 1996, 56:1050-1055). High efficiency gene transfer to subcutaneous tumours has been claimed with significant cell transfection
15 enhancement and better distribution characteristics over intra-tumoural injection procedures.

Liposomes work by surrounding hydrophilic molecules with hydrophobic molecules to facilitate cell entry. Liposomes are unilamellar or multilamellar spheres made from lipids. Lipid composition and manufacturing processes affect liposome structure. Other molecules can be incorporated into the lipid membranes. Liposomes can be anionic or cationic.
20 Nicolau *et al.*, Proc. Natl. Acad. Sci. U.S.A., 1983, 80: 1068-1072 has published on insulin expression from anionic liposomes injected into rats. Anionic liposomes mainly target the reticuloendothelial cells of the liver, unless otherwise targeted. Molecules can be incorporated into the surface of liposomes to alter their behavior, for example cell-selective delivery (Wu, G.Y. and Wu, C.H., J. Biol. Chem., 1987, 262: 4429-4432).

25 Felgner *et al.*, Proc. Nat. Acad. Sci. U.S.A., 1987, 84: 7413-7417 has published on cationic liposomes, demonstrated their binding of nucleic acids by electrostatic interactions and shown cell entry. Intravenous injection of cationic liposomes leads to transgene expression in most organs on injection into the afferent blood supply to the organ. Cationic liposomes can be administered by aerosol to target lung epithelium (Brigham *et al.*, Am. J.
30 Med. Sci., 1989, 298: 278-281). Patent references on liposomes are: WO 90/11092, WO 91/17424, WO 91/16024, WO 93/ 14788 (Vical) and; WO 90/01543 (Intracel).

In-Vivo studies with cationic liposome transgene delivery have been published by: Nabel *et al.*, Rev. Hum. Gene Ther., 1994, 5: 79-92 ; Hyde *et al.*, Nature, 1993, 362: 250-255 and ; Conary *et al.*, J. Clin. Invest., 1994, 93: 1834-1840).

Microparticles are being studied as systems for delivery of DNA to phagocytic cells
5 such approaches have been pursued by Pangaea Pharmaceuticals in their ENDOSHERE™ DNA microencapsulation delivery system which has been used to effect more efficient transduction of phagocytic cells such as macrophages which ingest the microspheres. The microspheres encapsulate plasmid DNA encoding potentially immunogenic peptides which when expressed lead to peptide display via MHC molecules on the cell surface which can
10 stimulate immune response against such peptides and protein sequences which contain the same epitopes. This approach is presently aimed towards a potential role in anti-tumour and pathogen vaccine development but may have other possible gene therapy applications.

In the same way as synthetic polymers have been used to package DNA natural viral coat proteins which are capable of homogeneous self-assembly into Virus-like particles
15 (VLPs) have been used to package DNA. The major structural coat protein VP1 of human polyoma virus can be expressed as a recombinant protein and is able to package plasmid DNA during self-assembly into a VLP. The resulting particles can be subsequently used to transduce various cell lines, while preliminary studies show little immunogenic response to such VP1 based VLPs. Such systems may offer an attractive intermediate between synthetic
20 polymer non-viral vectors and the alternative viral delivery systems since they may offer combined advantages e.g. simplicity of production and high level transduction efficiency.

To improve the specificity of gene delivery and expression the therapeutic gene the inclusion of targeting elements into the delivery vehicles and the use of regulatory expression elements have been investigated both singlulary and in combination in many of the previously
25 described delivery systems.

Improvements in DNA vectors have also been made and are likely applicable to all of the non-viral delivery systems. These include the use of supercoiled minicircles reported by RPR Gencell (which do not have bacterial origins of replication nor antibiotic resistance genes and thus are potentially safer as they exhibit a high level of biological containment), episomal
30 expression vectors as developed by Copernicus Gene Systems Inc (replicating episomal expression systems where the plasmid amplifies within the nucleus but outside the chromosome and thus avoids genome integration events) and T7 systems as developed by

Progenitor (a strictly a cytoplasmic expression vector in which the vector itself expresses phage T7 RNA polymerase and the therapeutic gene is driven from a second T7 promoter, using the polymerase generated by the first promoter). Other, more general improvements to DNA vector technology include use of cis-acting elements to effect high levels of expression
5 (Vical), sequences derived from alphoid repeat DNA to supply once-per-cell-cycle replication and nuclear targeting sequences (from EBNA-1 gene (Calos at Stanford, with Megabios); SV40 early promoter/enhancer or peptide sequences attached to the DNA).

Targeting systems based on cell receptor recognition by ligand linked to DNA have been described by Michael, S.I. and Curiel, D.T., *Gene Therapy*, 1994, 1: 223-232. Using the
10 ligand recognized by such a receptor the DNA becomes selectively bound and internalized into the target cell (Wu, G.Y. and Wu, C.H., *J. Biol. Chem.*, 1987, 262: 4429-4432). Poly-L-lysine (PLL), a polycation, has been used to couple a variety of protein ligands to DNA by chemical cross-linking methods. DNA is electrostatically bound to PLL-ligand molecules. Targetting systems have been published by Zenke *et al.*, *Proc. Nat. Acad. Sci. U.S.A.*, 1990,
15 87: 3655-3659 using transferrin receptor; Wu, G.Y. and Wu, C.H., *J. Biol. Chem.*, 1987, 262: 4429-4432 using the asialoorosomucoid receptor, and Batra *et al.*, *Gene Therapy*, 1994, 1: 255-260, using cell surface carbohydrates. Agents such as chloroquine or co-localised adenovirus can be used to reduce DNA degradation in the lysosomes (*see* Fisher, K.J. and Wilson, J.M., *Biochem. J.*, 1994, 299, 49-58). Cristiano *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*,
20 1993, 90: 11548-11552 has constructed adenovirus-DNA-ligand complexes. Patent references on receptor mediated endocytosis are: WO 92/05250 (asialoglycoproteins, University of Connecticut) and US 5354844 (transferrin receptor, Boehringer).

DNA and ligand can be coated over the surface of the adenovirus to create a coated adenovirus (Fisher, K.J. and Wilson, J.M., *Biochem. J.*, 1994, 299, 49-58). However the
25 presence of two receptor pathways for DNA entry (ligand receptor and adenovirus receptor) reduces the specificity of this delivery system but the adenovirus receptor pathway can be eliminated by using an antibody against adenovirus fiber protein as the means for linkage to DNA (Michael, S.I. and Curiel, D.T., *Gene Therapy*, 1994, 1: 223-232). Use of purified endosomalytic proteins rather than intact adenovirus particles is another option (Seth, P., *J.*
30 *Virol.*, 1994, 68: 1204-1206).

The expression of a gene construct of the invention at its target site is preferably under the control of a transcriptional regulatory sequence (TRS). A TRS is a promoter optionally

- 16 -

combined with an enhancer and/or an control element such as a genetic switch described below.

One example of a TRS is a "genetic switch" that may be employed to control expression of a gene construct of the invention once it has been delivered to a target cell.

5 Control of gene expression in higher eucaryotic cells by procaryotic regulatory elements (which are preferred for the present invention) has been reviewed by Gossen et al in TIBS, 18th December 1993, 471-475. Suitable systems include the *E.coli lac* operon and the especially preferred *E.coli tetracycline* resistance operon. References on the tetracycline system include Gossen et al (1995) Science 268, 1766; Damke et al (1995) Methods in

10 Enzymology 257, Academic Press; Yin et al (1996) Anal. Biochem. 235, 195 and; patents US 5464758, US 5589362, WO 96/01313 and WO 94/29442 (Bujard). An ecdysone based switch (International Patent Appln No. PCT/GB96/01195, Publication No. WO 96/37609, Zeneca) is another option. Other options are listed below. Connaught Laboratories (WO-93/20218) describe a synthetic inducible eukaryotic promoter comprising at least two different classes of

15 inducible elements. Rhone-Poulenc Rorer (WO 96/30512) describe a *tetracycline*-related application for a conditional gene expression system. Ariad (WO 94/18317) describes a protein dimerisation based system for which *in vivo* activity has been shown. Bert O'Malley of the Baylor College of Medicine (WO 93/23431, US 5364791, WO 97/10337) describes a molecular switch based on the use of a modified steroid receptor. The Whitehead Institute

20 have an NF-KB inducible gene expression system (WO 88/05083). Batelle Memorial have described a stress inducible promoter (European patent EP 263908).

Examples of TRSs which are independent of cell type include the following: cytomegalovirus promoter/ enhancer, SV40 promoter/ enhancer and retroviral long terminal repeat promoter/ enhancer. Examples of TRSs which are dependent on cell type (to give an

25 additional degree of targeting) include the following promoters: carcinoembryonic antigen (CEA) for targeting colorectal, lung and breast; alpha-foetoprotein (AFP) for targeting transformed hepatocytes; tyrosine hydroxylase, choline acetyl transferase or neurone specific enolase for targeting neuroblastomas; insulin for targeting pancreas and; glial fibro acidic protein for targeting glioblastomas. Some oncogenes may also be used which are selectively

30 expressed in some tumours e.g. HER-2/neu or c-erbB2 in breast and N-myc in neuroblastoma.

Accordingly, a preferred gene construct for use as a medicament is a construct comprising a transcriptional regulatory sequence which comprises a promoter and a control

- 17 -

element which is a genetic switch to control expression of the gene construct. A preferred genetic switch control element is regulated by presence of tetracycline or ecdysone. A preferred promoter is dependent on cell type and is selected from the following promoters: carcinoembryonic antigen (CEA); alpha-foetoprotein (AFP); tyrosine hydroxylase; choline acetyl transferase; neurone specific enolase; insulin; glial fibro acidic protein; HER-2/neu; c-erbB2; and N-myc. Preferably the gene construct for use as a medicament described herein is packaged within an adenovirus for delivery to the mammalian host. A general review of targeted gene therapy is given in Douglas *et al.*, Tumor Targeting, 1995, 1: 67-84.

The antibody encoded by the gene construct of the invention may be any form of antibody construct such as for example F(ab')₂; F(ab'), Fab, Fv, single chain Fv & V-min. Any suitable antibody construct is contemplated, for example a recently described antibody fragment is "L-F(ab)₂" as described by Zapata (1995) in Protein Engineering, 8, 1057-1062. Disulphide bonded Fvs are also contemplated. For constructs based on CPG2 enzyme, Fab fragment constructs dimerised through enzyme dimerisation are preferred. Non-human antibodies may be humanised for use in humans to reduce host immune responses. A humanized antibody, related fragment or antibody binding structure is a polypeptide composed largely of a structural framework of human derived immunoglobulin sequences supporting non human derived amino acid sequences in and around the antigen binding site (complementarity determining regions or CDRs). Appropriate methodology has been described for example in detail in WO 91/09967, EP 0328404 and Queen et al. Proc Natl Acad Sci 86,10029, Mountain and Adair (1989) Biotechnology and Genetic Engineering Reviews 10, 1 (1992) although alternative methods of humanisation are also contemplated such as antibody veneering of surface residues (EP 519596, Merck/NIH, Padlan et al).

According to another aspect of the present invention there is provided a matched two component system designed for use in a mammalian host in which the components comprise:

- (i) a first component that comprises a gene construct encoding a cell targeting antibody and a heterologous prodrug activating enzyme wherein the gene construct is capable of expressing the antibody and enzyme as a conjugate within a target cell in the mammalian host and wherein the conjugate can leave the cell thereafter for selective localisation at a cell surface antigen recognised by the antibody and;
- (ii) a second component that comprises a prodrug which can be converted into an active drug by the enzyme.

- 18 -

Antibody directed enzyme prodrug therapy (ADEPT) is a known cancer therapeutic approach. ADEPT uses a tumour selective antibody conjugated to an enzyme. The conjugate is administered to the patient (usually intravenously), allowed to localise at the tumour site(s) and clear from the blood and other normal tissues. A prodrug is then administered to the patient which is converted by the enzyme (localised at the tumour site) into a cytotoxic drug which kills the tumour cells.

The present invention can be applied to any ADEPT system. Suitable examples of ADEPT systems include those based on any of the following enzymes: carboxypeptidase G2; carboxypeptidase A; aminopeptidase; alkaline phosphatase; glycosidases; β -glucuronidase; penicillin amidase; β -lactamase; cytosine deaminase; nitroreductase; or mutant host enzymes including carboxypeptidase A, carboxypeptidase B, and ribonuclease. Suitable references on ADEPT systems include Melton RG (1996) in J. National Cancer Institute 88, 1; Niculescu-Duvaz I (1995) in Current Medicinal Chemistry 2, 687; Knox RJ (1995) in Clin. Immunother. 3, 136; WO 88/07378 (CRCT); Blakey et al, Cancer Res. 56, 3287-92, 1996; US 5587161 (CRCT and Zeneca); WO 97/07769 (Zeneca); and WO 95/13095 (Wellcome). The heterologous enzyme may be in the form of a catalytic antibody; see for example EP 745673 (Zeneca). A review articles on ADEPT systems include Hay & Denny (1996), Drugs of the Future, 21(9), 917-931 and Blakey (1997), Exp. Opin. Ther. Patents, 2(9), 965-977.

A preferred matched two component system is one in which:

the first component comprises a gene encoding the heterologous enzyme CPG2; and the second component prodrug is selected from N-(4-[N,N-bis(2-iodoethyl)amino]-phenoxy-carbonyl)-L-glutamic acid, N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic- γ -(3,5-dicarboxy)anilide or N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic acid or a pharmaceutically acceptable salt thereof. Preferred prodrugs for use with CPG2 are described in the following US patents from Zeneca Limited and Cancer Research Campaign Technology Limited: US 5714148, US 5405990, 5587161 & 5660829.

In another aspect of the invention there is provided a method for the delivery of a cytotoxic drug to a site which comprises administering to a host a first component that comprises a gene construct as defined herein; followed by administration to the host of a second component that comprises a prodrug which can be converted into a cytotoxic drug by the heterologous enzyme encoded by the first component. A preferred method for delivery of

- 19 -

a cytotoxic drug to a site is one in which the first component comprises a gene encoding the heterologous enzyme CPG2; and the second component prodrug is selected from N-(4-[N,N-bis(2-iodoethyl)amino]phenoxy-carbonyl)-L-glutamic acid, N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic-gamma-(3,5-dicarboxy)anilide or N-5 (4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic acid or a pharmaceutically acceptable salt thereof.

Abbreviations used herein include:

AAV	Adeno-associated virus
ADEPT	antibody directed enzyme prodrug therapy
AFP	alpha-foetoprotein
AMIRACS	Antimetabolite with Inactivation of Rescue Agents at Cancer Sites
APS	ammonium persulfate
b.p.	base pair
BPB	bromophenol blue
CDRs	complementarity determining regions
CEA	Carcinoma Embryonic Antigen
CL	constant domain of antibody light chain
CPB	carboxypeptidase B
CPG2	carboxypeptidase G2
CPG2 R6	carboxypeptidase G2 mutated to prevent glycosylation on expression in eucaryotic cells, see Example 1d
DAB	substrate 3,3'-diaminobenzidine tetrahydrochloride
DEPC	diethylpyrocarbonate
DMEM	Dulbecco's modified Eagle's medium
ECACC	European Collection of Animal Cell Cultures
EIA	enzyme immunoassay
ELISA	enzyme linked immunosorbent assay
FAS	folinic acid supplemented
FCS	foetal calf serum

- 20 -

Fd	heavy chain of Fab, Fab' or F(ab') ₂ optionally containing a hinge
GDEPT	gene directed enzyme prodrug therapy
HAMA	Human Anti Mouse Antibody
HCPB	human carboxypeptidase B, preferably pancreatic
hinge (of an IgG)	a short proline rich peptide which contains the cysteines that bridge the 2 heavy chains
HRPO or HRP	horse radish peroxidase
IRES	internal ribosome entry site
MTX	methotrexate
NCA	non-specific cross reacting antigen
NCIMB	National Collections of Industrial and Marine Bacteria
OPD	<i>ortho</i> -phenylenediamine
PBS	phosphate buffered saline
PCR	polymerase chain reaction
PGP	<u>N</u> -(4-[<u>N,N</u> -bis(2-chloroethyl)amino]-phenoxy carbonyl)- <u>L</u> -glutamic acid
preproCPB	proCPB with an N-terminal leader sequence
proCPB	CPB with its N-terminal pro domain
scFv	single chain Fv
SDS-PAGE	sodium dodecyl sulphate - polyacrylamide gel electrophoresis
SSC	salt sodium citrate
TBS	Tris-buffered Saline
Temed	N,N,N',N'-tetramethylethylenediamine
TFA	trifluoroacetic acid
TRS	transcriptional regulatory sequence
VDEPT	virus-directed enzyme prodrug therapy
VH	variable region of the heavy antibody chain
VK	variable region of the light antibody chain

In this specification conservative amino acid analogues of specific amino acid sequences are contemplated which retain the relevant biological properties of the component of the invention but differ in sequence by one or more conservative amino acid substitutions, deletions or additions. However the specifically listed amino acid sequences are preferred. Typical conservative amino acid substitutions are tabulated below.

Original	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Lys; Arg	Gln
Asp (D)	Glu	Glu
Cys (C)	Ser	Ser
Gln (Q)	Asn	Asn
Glu (E)	Asp	Asp
Gly (G)	Pro	Pro
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Leu; Val; Ile; Ala	Leu
Pro (P)	Gly	Gly
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe

- 22 -

Val (V) Ile; Leu; Met; Phe; Leu
 Ala; Norleucine

Amino acid nomenclature is set out below.

Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartic Acid	Asp	D
Cysteine	Cys	C
Glutamic Acid	Glu	E
Glutamine	Gln	Q
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V
Any Amino Acid	Xaa	X

5

In this specification nucleic acid variations (deletions, substitutions and additions) of specific nucleic acid sequences are contemplated which retain which the ability to hybridise under stringent conditions to the specific sequence in question. Stringent conditions are defined as 6xSSC, 0.1 % SDS at 60° for 5 minutes. However specifically listed nucleic acid

sequences are preferred. It is contemplated that chemical analogues of natural nucleic acid structures such as "peptide nucleic acid" (PNA) may be an acceptable equivalent, particularly for purposes that do not require translation into protein (Wittung (1994) Nature 368, 561).

The invention will now be illustrated by reference to the following non-limiting

5 Examples. Temperatures are in degrees Celsius.

Figure 1 shows a representation of the fusion gene construct comprising A5B7 antibody heavy chain Fd fragment linked at its C-terminus via a flexible (G4S)₃ peptide linker to the N-terminus of CPG2 polypeptide. SS represents the signal sequence. L represents a linker sequence. CPG2/R6 represents CPG2 with its glycosylation sites nullified through mutation
10 as explained in the text.

Figure 2a shows a representation of (Fab-CPG2)₂ fusion protein with dimerisation taking place through non-covalent bonding between two CPG2 molecules.

Figure 2b shows a representation of a F(ab')₂ antibody fragment.

Figure 3 shows a cell based ELISA assay of secreted fusion protein material. Only the CEA
15 positive line has increased levels of binding with increasing amounts of added fusion protein whereas the CEA negative cell line has only constant background binding levels throughout. The vertical axis represents optical density readings measured at 490 nm and the horizontal axis the amount of added fusion protein measured in ng of protein. The graph shows data obtained from an experiment where a number of cell lines and a negative control (no
20 cells) were incubated with increasing amounts of fusion protein using the cell assay described in Example 6. The results show that only the LoVo (CEA positive) cell line showed an increasing OD490 reading corresponding to increasing amounts of added fusion protein. All other cell lines (CEA negative) and the control (no cells) showed only a background OD490 nm reading which did not increase with the addition of fusion protein. These results provide
25 evidence that the fusion protein material binds specifically to a CEA positive cell line in a dose dependant manner and do not bind to CEA negative lines.

Figure 4 shows retention of secreted fusion protein to recombinant LoVo tumour cells. The vertical axis represents optical density readings measured at 490 nm and the horizontal axis the amount of added anti-CEA antibody (IIE6) measured in ng/ml of protein. The experiment
30 was performed as described in Example 7 using three different cell lines, recombinant LoVo and Colo320DM lines (which themselves secrete fusion protein) and a control parental LoVo line which does not secrete fusion protein. Firstly, the cell lines were fixed, and washed to

- 24 -

remove the existing supernatant and any unbound material after which increasing concentrations of the anti-CEA antibody (IIE6) were added to the fixed cells. The assay was developed as described in the text to determine the level of retention of any secreted material and whether further added antibody would increase the signal. The results showed that

5 without added anti-CEA antibody the control parental Lovo line exhibited only a background OD490 nm reading (as expected) whereas the recombinant LoVo line gave a very strong OD 490 nm reading indicating that the fusion protein material was being retained on the CEA positive LoVo cells. The CEA negative recombinant Colo320DM gave a much weaker reading than the LoVo cells but the signal was higher than background (possibly due

10 to none fixing of the secreted antibody early in the assay method). Increasing concentrations of the anti-CEA antibody (IIE6) added to the fixed cells showed a dose related response in the case of the parental LoVo cells thus indicating that they are CEA positive and can bind CEA binding material (such as the fusion protein if present or added). The recombinant Colo320DM and LoVo cells showed little increase in overall OD490 signal with increasing

15 amounts of added antibody with the exception of the LoVo cells which appear to show a slight response at the highest antibody dose. Since the recombinant Colo320DM are CEA negative no increase in signal due to anti-CEA antibody the results for these cells would be expected. In the case of the recombinant LoVo cells the addition signal due the amounts of antibody added in this assay may be swamped except at the highest dose due to the relative

20 strength of the original signal.

Figure 5 shows retention of secreted fusion protein to recombinant LoVo tumour cells. The vertical axis represents median tumour volume (cm^3) and the horizontal axis time in day after dosing of the prodrug. The experiment was performed as described in Example 12 using 60 mg/kg doses of prodrug. The results show that the control GAD(c) (none prodrug treated)

25 tumours grew to 6 times their initial size by 11 days (post-dose day) at which time the tumours were harvested. The prodrug treated tumours GAD(d) show a significantly slower growth rate and by day 16 (post-dose day) have only reached 3 times their initial size. This data indicates at least an 11 day tumour growth delay.

In the Examples below, unless otherwise stated, the following methodology and

30 materials have been applied.

DNA is recovered and purified by use of GENECLEAN™ II kit (Strattech Scientific Ltd. or Bio 101 Inc.). The kit contains: 1) 6M sodium iodide; 2) a concentrated solution of

sodium chloride, Tris and EDTA for making a sodium chloride/ethanol/water wash; 3) Glassmilk- a 1.5 ml vial containing 1.25 ml of a suspension of a specially formulated silica matrix in water. This is a technique for DNA purification based on the method of Vogelstein and Gillespie published in Proceedings of the National Academy of Sciences USA (1979) Vol 5 76, p 615. Briefly, the kit procedure is as follows. To 1 volume of gel slice is added 3 volumes of sodium iodide solution from the kit. The agarose is melted by heating the mix at 55° for 10 min then Glassmilk (5-10 ml) is added, mixed well and left to stand for 10 min at ambient temperature. The glassmilk is spun down and washed 3 times with NEW WASH™ (0.5 ml) from the kit. The wash buffer is removed from the Glassmilk and DNA is eluted by 10 incubating the Glassmilk with water (5-10 ml) at 55° for 5-10 min. The aqueous supernatant containing the eluted DNA is recovered by centrifugation. The elution step can be repeated and supernatants pooled.

Competent E. coli DH5α cells were obtained from Life Technologies Ltd (MAX™ efficiency DH5α competent cells).

15 Mini-preparations of double stranded plasmid DNA were made using the RPM™ DNA preparation kit from Bio101 Inc. (cat. No 2070-400) or a similar product - the kit contains alkaline lysis solution to liberate plasmid DNA from bacterial cells and glassmilk in a spinfilter to adsorb liberated DNA which is then eluted with sterile water or 10mM Tris-HCl, 1mM EDTA, pH 7.5.

20 The standard PCR reaction contains 100 ng of plasmid DNA (except where stated), 5 µl dNTPs (2.5 mM), 5 µl 10x Enzyme buffer (500 mM KCl, 100 mM Tris pH 8.3), 15mM MgCl₂ and 0.1 % gelatin), 1 µl of a 25 pM/ µl stock solution of each primer, 0.5 µl thermostable DNA polymerase and water to obtain a volume of 50 µl. Standard PCR conditions were: 15 cycles of PCR at 94° for 90 s; 55° for 60 s; 72° for 120 s, ending the last 25 cycle with a further 72° for 10 min incubation.

AMPLITAQ™, available from Perkin-Elmer Cetus, is used as the source of thermostable DNA polymerase.

General molecular biology procedures can be followed from any of the methods described in "Molecular Cloning - A Laboratory Manual" Second Edition, Sambrook, Fritsch 30 and Maniatis (Cold Spring Harbor Laboratory, 1989).

Serum free medium is OPTIMEM™ I Reduced Serum Medium, GibcoBRL Cat. No.

- 26 -

31985. This is a modification of Eagle's Minimum Essential Medium buffered with Hepes and sodium bicarbonate, supplemented with hypoxanthine, thymidine, sodium pyruvate, L-glutamine, trace elements and growth factors.

LIPOFECTIN™ Reagent (GibcoBRL Cat. No. 18292-011) is a 1:1 (w/w) liposome formulation of the cationic lipid N-[1-(2,3-dioleoyloxy)propyl]-n,n,n-trimethylammonium chloride (DOTMA) and dioleoyl phosphatidylethanolamine (DOPE) in membrane filtered water. It binds spontaneously with DNA to form a lipid-DNA complex - see Felgner *et al.* in Proc. Natl. Acad. Sci. USA (1987) 84, 7431.

G418 (sulphate) is GENETICIN™, GibcoBRL Cat. No 11811, an aminoglycoside antibiotic related to gentamicin used as a selecting agent in molecular genetic experiments;

For the CEA ELISA each well of a 96 well immunoplate (NUNC MAXISORB™) was coated with 50ng CEA in 50 mM carbonate/bicarbonate coating buffer pH9.6 (buffer capsules - Sigma C3041) and incubated at 4° overnight. The plate was washed three times with PBS-TWEENT™ (PBS + 0.05 % TWEENT™ 20) and then blocked 150 µl per well of 1 % BSA in PBS-TWEENT™ for 1 hour at room temperature. The plate was washed three times with PBS-TWEENT™, 100 µl of test sample added per well and incubated at room temperature for 2 hours. The plate was washed three times with PBS-TWEENT™, 100 µl per well of a 1/500 dilution of HRPO-labelled goat anti-human kappa antibody (Sigma A 7164) was added in 1 % BSA in PBS-TWEENT™ and incubated at room temperature on a rocking platform for at least 1 hour. The plate was washed three times with PBS-TWEENT™ and then once more with PBS. To detect binding, add 100µl per well of developing solution (one capsule of phosphate-citrate buffer - Sigma P4922 - dissolved in 100 ml H₂O to which is added one 30 mg tablet o-phenylenediamine dihydrochloride - Sigma P8412) and incubated for up to 15 minutes. The reaction was stopped by adding 75 µl 2M H₂SO₄, and absorbance read at 490 nm.

The CEA ELISA using an anti CPG2 reporter antibody was essentially as above but instead of HRPO-labelled goat anti-human kappa antibody an 1/1000 dil. of a rabbit anti-CPG2 polyclonal sera was added, in 1 % BSA in PBS-TWEENT™ and incubated at room temperature on a rocking platform for at 2 hours. The plate was washed three times with PBS-TWEENT™. A 1/2000 dilution of a goat anti-rabbit HRPO labelled antibody (Sigma A-6154) was then added and incubated at room temperature on a rocking platform for 1 hour, the

- 27 -

plate was washed three times with PBS-TWEEN™ and once with PBS. To detect binding add 100µl per well developing solution (one capsule of phosphate-citrate buffer - Sigma P4922 - dissolved in 100 ml H₂O to which is added one 30 mg tablet *o*-phenylenediamine dihydrochloride - Sigma P8412) and incubated for up to 15 minutes. The reaction was stopped by adding 75µl 2M H₂SO₄, and absorbance read at 490nm.

Western blot analysis of transfection supernatants was performed as follows.

- 10 % mini gels** for analysis of fusion protein transfections were prepared using a suitable mini gel system (HOEFER MIGHTY SMALL™). 10 % running gel is: 20 ml acrylamide, 6 ml 10 x running gel buffer; 34 ml H₂O; 300 ml 20 % SDS; 600 µl APS; 30 µl Temed.
- 10 Running gel buffer 10x is 3.75 M Tris pH 8.6. 6 % stacking gel is: 9 ml acrylamide; 4.5 ml 10x stacking gel buffer; 31.5 ml H₂O; 225 µl 20 % SDS 450 µl 10 % APS; 24 µl Temed). Stacking gel buffer 10x is 1.25 M Tris pH 6.8. Electrophoresis buffer 5x for SDS/PAGE is 249 mM Tris, 799 mM glycine, 0.6 % w/v SDS (pH not adjusted).

- Preparation of samples** 2 x Laemmli buffer is 0.125 M Tris; 4 % SDS; 30 % glycerol; 4 M urea; 0.002 % BPB optionally containing 5 % β-mercaptoethanol.
- Supernatants:** 25 µl sample + 25 µl 2 x Laemmli buffer; 40 µl loaded. **Standards F(ab')₂ and CPG2:** 2 µl of 10 ng/ml of standard; 8 µl of H₂O; 10 µl 2x Laemmli buffer (-mercaptoethanol); 20 µl loaded. **Molecular weight markers** (Amersham RAINBOW™): 8 µl sample; 8 µl 2x Laemmli buffer (+ mercaptoethanol); 16 µl loaded. **Running conditions:** 20 30 milliamps until dye front at bottom of gel (approx. 1 hour). **Blotting:** using a semi dry blotter (LKB) onto nitrocellulose membrane. Milliamps = 0.7 x cm², for 45 minutes. **Blocking:** 5 % dried skimmed milk in PBS-TWEEN™ for 40 minutes.

Detection of F(ab')₂: goat anti human kappa light chain HRPO labelled antibody, 1/2500 in 0.5 % dried skimmed milk in PBS-TWEEN™ incubated overnight.

- 25 **Detection of CPG2:** mouse anti-CPG2 monoclonal (1/2000 in 0.5 % dried skimmed milk in PBS-TWEEN™ incubated overnight; goat anti mouse kappa light chain HRPO labelled antibody -Sigma 674301- (1/10000 in 0.5 % dried skimmed milk in PBS-TWEEN™) incubated for at least 2 hours.

- Development of Blot:** Chemiluminescence detection of HRPO based on luminol substrate in the presence of enhancer was used (Pierce SUPERSIGNAL™ Substrate). Substrate working solution was prepared as follows: recommended volume: 0.125 ml/cm² of

- 28 -

blot surface. Mix equal volumes of luminol/enhancer solution and stable peroxide solution, incubate blot with working solution for 5-10 minutes, remove solution and place blot in a membrane protector and expose against autoradiographic film (usually between 30 seconds and 5 minutes).

5 **Microorganism deposits:** Plasmid pNG3-Vkss-HuCk was deposited at The National Collections of Industrial and Marine Bacteria (NCIMB), 23 St Machar Drive, Aberdeen AB2 1RY, Scotland, United Kingdom on 11-April-1996 under deposit reference number NCIMB 40798 in accordance with the Budapest Treaty. Plasmid pNG4-VHss-HuIgG2CH1' was deposited at The National Collections of Industrial and Marine Bacteria (NCIMB), 23 St Machar Drive, Aberdeen AB2 1RY, Scotland, United Kingdom on 11-April-1996 under deposit reference number NCIMB 40797 in accordance with the Budapest Treaty. Plasmid pNG3-Vkss-HuCk-NEO was deposited at The National Collections of Industrial and Marine Bacteria (NCIMB), 23 St Machar Drive, Aberdeen AB2 1RY, Scotland, United Kingdom on 11-April-1996 under deposit reference number NCIMB 40799 in accordance with the Budapest Treaty. Plasmid **pICI266** was deposited under accession number NCIMB 40589 on 11Oct93 under the Budapest Treaty at the National Collections of Industrial and Marine Bacteria Limited (NCIMB), 23 St. Machar Drive, Aberdeen, AB2 1RY, Scotland, U.K.

Trypsinisation: Trypsin EDTA (Gibco BRL 45300-019) and Hanks balanced salt solution (HBSS; Gibco BRL 14170-088) were pre-warmed in a 37° waterbath. Existing media was removed from cultures and replaced with a volume of HBSS (which is half the previous media volume) and the layer of cells washed by carefully rocking the plate or flask so as to remove any residual serum containing media. The HBSS was removed and a volume of Trypsin solution (which is one quarter of the original media volume) added, with gently rocking the flask to ensure the cell layer was completely covered and left for 5 min. Trypsin was inactivated by addition of of the appropriate normal culture media (2x the volume of the trypsin solution). The cell suspension was then either cell counted or further diluted for continued culture depending on the procedure to be performed.

Heat Inactivation of Foetal Calf Serum (FCS): FCS (Viralex A15-651 accredited batch - Non European) was stored at -20°. For use, the serum was completely thawed at 4° overnight. The next day, the serum was incubated for 15 min in a 37° waterbath and then transferred to a 56° waterbath for 15 min. The serum was removed and allowed to cool to room temperature before it was split in to 50 ml aliquots and stored at -20°C

- 29 -

Normal DMEM Media (using Gibco BRL components): To 500 ml DMEM (41966-086) add 12.5ml Hepes (15630-056); 5ml NEAA (11140-035); 5 ml pen/strep (10378-016); and 50 ml heat inactivated FCS.

FAS Media (using Gibco BRL components unless stated otherwise): 490 ml DMEM 5 (41966-086); 12.5 ml Hepes (15630-056); 5ml non-essential amino acids (11140-035); 5 ml pen/strep (10378-016); 5 ml vitamins (11120-037); 5ml basal amino acids (51051-019); Folinic Acid (Sigma F8259) to a final media concentration of 10 µg/ml ; 50 ml heat inactivated FCS; 5 ml dNTP mix; and G418 50 mg/ml stock solution (to produce the appropriate selection concentration).

10 **dNTP mix:** 35mg G (Sigma G6264), 35mg C (Sigma C4654), 35mg A (Sigma A4036), 35mg U (SigmaU3003), 125mg T (Sigma T1895) were dissolved in 100ml water, filter sterilised, and stored at -20°.

G418 Selection: for LoVo cells (ATCC CCL 229) selection was performed at 1.25 mg/ml, for HCT116 (ATCC CCL 247) cells and for Colo320DM (ATCC CCL 220) cells
15 selection was performed at 1.5 mg/ml unless stated otherwise.

BLUESCRIPT™ vectors were obtained from Stratagene Cloning Systems.

Tet-On gene expression vectors were obtained from Clontech (Palo Alto, California) cat. no. K1621-1.

Unless stated otherwise or apparent from the context used, antibody-CPG2 fusion
20 constructs referred to in the Examples use mutated CPG2 to prevent glycosylation.

Example 1

Construction of an (A5B7 Fab-CPG2)₂ fusion protein

The construction of a (A5B7 Fab-CPG2)₂ enzyme fusion was planned with the aim of
25 obtaining a bivalent human carcinoembryonic antigen (CEA) binding molecule which also exhibits CPG2 enzyme activity. To this end the initial construct was designed to contain an A5B7 antibody heavy chain Fd fragment linked at its C-terminus via a flexible (G4S)₃ peptide linker to the N-terminus of the CPG2 polypeptide (Figure 1).

The antibody A5B7 binds to human carcinoembryonic antigen (CEA) and is
30 particularly suitable for targeting colorectal carcinoma or other CEA antigen bearing cells (the importance of CEA as a cancer associated antigen is reviewed by Shively, J.E. and Beatty, J.D. in "CRC Critical Reviews in Oncology/Hematology", vol 2, p355-399, 1994).

- 30 -

The CPG2 enzyme is naturally dimeric in nature, consisting of two associated identical polypeptide subunits. Each subunit of this molecular dimer consists of a larger catalytic domain and a second smaller domain that forms the dimer interface.

In general, antibody (or antibody fragment)-enzyme conjugate or fusion proteins should be at least divalent, that is to say capable of binding at least 2 tumour associated antigens (which may be the same or different). In the case of the (A5B7 Fab-CPG2)₂ fusion protein, dimerisation of the enzyme component takes place after expression, as with the native enzyme, thus forming an enzymatic molecule which contains two Fab antibody fragments (and is thus bivalent with respect to antibody binding sites) and two molecules of CPG2 (Figure 2a).

a) Cloning of the A5B7 antibody genes

Methods for the preparation, purification and characterisation of recombinant murine A5B7 F(ab')₂ antibody have been published (International Patent Application, Zeneca Limited, WO 96/20011, see Reference Example 5 therein). In Reference Example 5, section f thereof, the A5B7 antibody genes were cloned into vectors of the GS-SYSTEM™ (Celltech), see International Patent Applications WO 87/04462, WO 89/01036, WO 86/05807 and WO 89/10404, with the A5B7 Fd cloned into pEE6 and the light chain into pEE12. These vectors were the source of the A5B7 antibody genes for the construction of the A5B7 Fab-CPG2 fusion protein.

20 b) Chimaeric A5B7 vector constructs

The A5B7 murine antibody variable regions were amplified by PCR from the pEE6 and pEE12 plasmid vectors using appropriate PCR primers which included the necessary restriction sites for direct in frame cloning of the heavy and light chain variable regions into the vectors pNG4-VHss-HuIgG2CH1' (NCIMB deposit no. 40797) and pNG3-Vkss-HuCK-NEO (NCIMB deposit no. 40799) respectively. The resulting vectors were designated pNG4/A5B7VH-IgG2CH1' (A5B7 chimaeric heavy chain Fd') and pNG3/A5B7VK-HuCK-NEO (A5B7 chimaeric light chain).

c) Cloning of the CPG2 gene

The CPG2 coding gene may be obtained from Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom. CPG2 may also be obtained by recombinant techniques. The nucleotide coding sequence for CPG2 has been published by Minton, N.P. *et al.*, Gene, (1984) 31, 31-38. Expression of the coding sequence

has been reported in *E. coli* (Chambers, S.P. et al., Appl. Microbiol, Biotechnol. (1988), 29, 572-578) and in *Saccharomyces cerevisiae* (Clarke, L. E. et al., J. Gen Microbiol, (1985) 131, 897-904). In addition the CPG2 gene may be produced as a synthetic DNA construct by a variety of methods and used as a source for further experiments. Total gene synthesis has been described by M. Edwards in Am. Biotech. Lab (1987), 5, 38-44, Jayaraman et al. (1991) Proc. Natl. Acad. Sci. USA 88, 4084-4088, Foguet and Lubbert (1992) Biotechniques 13, 674-675 and Pierce (1994) Biotechniques 16, 708.

In preparation for the cloning the CPG2 gene the vector pNG3-Vkss was constructed which is a simple derivative of pNG3-Vkss-HuCk-NEO (NCIMB deposit no. 40799). This vector was constructed by first removing the Neomycin gene (since it contained an EcoRI restriction enzyme site) by digestion with the restriction enzyme XbaI, after which the vector fragment was isolated and then religated to form the plasmid pNG3/Vkss-HuCk. This intermediate vector was digested with the enzymes SacII and EcoRI, which excised the HuCk gene fragment. The digest was then loaded on a 1 % agarose gel and the excised fragment separated from the remaining vector after which the vector DNA was cut from the gel and purified. Two oligonucleotides CME 00261 and CME 00262 (SEQ ID NO: 1 and 2) were designed and synthesised. These two oligonucleotides were hybridised by adding 200 pmoles of each oligonucleotide into a total of 30 µl of H₂O, heating to 95° and allowing the solution to cool slowly to 30°. 100 pmoles of the annealed DNA product was then ligated directly into the previously prepared vector and the ligation mix transformed into *E. coli*. In the clones obtained, the introduction of the DNA "cassette" produced a new polylinker sequence in preparation for the subsequent CPG2 gene cloning to produce the vector pNG3-Vkss.

The CPG2 structural gene encoding amino acid residues Q26-K415 inclusive was amplified by PCR using appropriate DNA oligonucleotide primers and standard PCR reaction conditions. The reaction product was analysed using a 1 % agarose gel, a band of the expected size (approximately 12000 b.p.) was excised, purified and eluted in 20µl H₂O. This material was then digested using the restriction enzyme SacII, after which the reaction was loaded on a 1 % agarose gel and a band of the expected size (approximately 250 b.p.) was excised and subsequently purified. This fragment was ligated into the plasmid vector pNG3VKss, which had been previously digested with the restriction enzyme SacII, dephosphorylated, run on a 1 % agarose gel, the linearised vector band excised, purified, and the ligation mix transformed

into *E.coli*. The resultant clones were analysed for the presence and orientation of the CPG2 SacII fragment by DNA restriction analysis using the enzymes BglII and FseI. Clones which appeared to have a fragment of the correct size and orientation were confirmed by DNA sequencing. This intermediate plasmid was called **pNG3-Vkss-SacIICPG2frag**. This
5 plasmid was digested with the restriction enzymes by AgeI and EcoRI, dephosphorylated and the vector fragment isolated. The original CPG2 gene PCR product was also digested with AgeI and EcoRI, an approximately 1000 bp. fragment isolated, ligated and transformed into *E.coli*. The resulting clones were analysed for a full length CPG2 gene (approximately 1200 bp.) by digestion with the restriction enzymes HindIII and EcoRI; clones with the correct size
10 insert were sequenced to confirm identity. Finally, this plasmid (pNG3/Vkss-CPG2) was digested with XbaI, dephosphorylated, a vector fragment isolated and the XbaI Neomycin gene fragment (approximately 1000 bp. which had also been isolated in the earlier stages) religated into the plasmid and transformed into *E.coli*. Resulting clones were checked for the presence and orientation of the Neomycin gene by individual digests with the enzymes XbaI
15 and EcoRI. This vector was called **pNG3-Vkss-CPG2-NEO**.

d) Construction of the CPG2 R6 variant

The plasmid pNG3-Vkss/CPG2-NEO was used as a template for the PCR mutagenesis of the CPG2 gene in order to mutate 3 potential glycosylation sites which had been identified within the natural bacterial enzyme sequence. The putative amino acid glycosylation sites (N-
20 X-T/S) were observed at positions 222 (N-I-T), 264 (N-W-T), and 272 (N-V-S) using the positional numbering published by Minton, N.P. et al., in *Gene*, (1984) 31, 31-38. The asparagine residue (N) of the 3 glycosylation sites was mutated to glutamine (Q) thus negating the glycosylation sites to avoid any glycosylation events affecting CPG2 expression or enzyme activity.

25 A PCR mutagenesis technique in which all 3 sites were mutated in a single reaction series was used to create the CPG2 R6 gene variant. The vector pNG3/Vkss/CPG2-NEO was used as the template for three initial PCR reactions. Reaction R1 used synthetic oligonucleotide sequence primers CME 00395 and CME 00397 (SEQ ID NOS: 3 and 4), reaction R2 used synthetic oligonucleotide sequence primers CME 00395 and CME 00399
30 (SEQ ID NOS: 3 and 5) and reaction R3 used synthetic oligonucleotide sequence primers CME 00396 and CME 00400 (SEQ ID NOS: 6 and 7). The products of PCR reactions R1 and R2 contained the mutated 222 and 264 + 272 glycosylation sites respectively, with the R3

- 33 -

product being a copy of the C-terminal segment of the CPG2 gene. The R2 and R3 products (R2 approximately 750 bp; R3 approximately 360 bp), after agarose gel separation and purification, were joined in a further PCR reaction. Mixtures of varying amounts of the products R2 and R3 were made and PCR reactions performed using the synthetic
5 oligonucleotides CME 00395 and CME 00396 (SEQ ID NOS: 3 and 6). The resulting product R4 (approximately 1200bps) was again PCR amplified using the oligonucleotides CME 00398 and CME 00396 (SEQ ID NOS: 8 and 6). The resulting product R5 (approximately 600 bp.) was joined to product R1 (approximately 620 b.p.) in a final PCR reaction performed using the oligonucleotides CME 00395 and CME 00396 (SEQ ID NOS:
10 3 and 6). The resulting PCR product R6 (approximately 1200 bp), which now contained all three mutated glycosylation sites, could be cloned (after digestion with the restriction enzymes AgeI and BsrGI and isolation of the resultant fragment) into the vector pNG3/Vkss-CPG2-Neo.(which had been previously cut with the restriction enzymes AgeI and Bsr GI and subsequently isolated). This created the desired DNA (SEQ ID NO: 9) encoding CPG2/R6
15 protein sequence (SEQ ID NO: 10) within the expression vector pNG3/Vkss-CPG2 R6-NEO.

e) Construction of the A5B7 heavy chain Fd-CPG2 fusion protein gene

The heavy chain antibody fragment and the CPG2 enzyme genes were both obtained by PCR amplification of plasmid templates. The plasmid pNG4/A5B7VH-IgG2CH1' was
20 amplified with primers CME 00966 (SEQ ID NO: 11) and CME 00969 (SEQ ID NO: 12) to obtain the A5B7 Fd component (approximately 300 b.p.) and the plasmid pNG3/Vkss/CPG2 R6-NEO was amplified with primers CME 00967 (SEQ ID NO: 13) and CME 00968 (SEQ ID NO: 14) to obtain the enzyme component (approximately 1350 b.p.). In each case the PCR reaction product was loaded and separated on a 1 % agarose gel, a band of the correct product
25 size excised, subsequently purified and eluted in 20 µl H₂O.

A further PCR reaction was performed to join (or splice) the two purified PCR reaction products together. Standard PCR reaction conditions were used with varying amounts (between 0.5 to 2 µl) of each PCR product but utilising 25 cycles (instead of the usual 15 cycles). The reaction product was analysed using a 1 % agarose gel and a band of
30 the expected size (approximately 1650 b.p.) was excised, purified and eluted in 20 µl H₂O. This material was then digested using restriction enzymes NheI and BamHI, after which a

- 34 -

band of the expected size (approximately 1600 b.p.) was recovered and purified. The vector pNG4/A5B7VH-IgG2CH1' was prepared to receive the above PCR product by digestion with restriction enzymes NheI and BamHI, after which the DNA was dephosphorylated and the larger vector band was separated from the smaller NheI/Bam HI fragment. The vector band was recovered, purified and subsequently the similarly restricted PCR product was ligated in to the prepared vector and the ligation mix transformed into *E. coli*. DNA was prepared from the clones obtained and subsequently sequenced to confirm the fusion gene sequence. A number of the clones were found to be correct and one of these clones (designated R2.8) was re-named pNG4/A5B7VH-IgG2CH1/CPG2 R6 (SEQ ID NO: 15 and SEQ ID NO: 16).

10 f) Co-transfection, transient expression

The plasmids pNG4/A5B7VH-IgG2CH1/CPG2 R6 (encoding the antibody chimaeric Fd-CPG2 fusion protein) and pNG3/A5B7VK-HuCK-NEO (encoding the antibody chimaeric light chain; SEQ ID NO: 17 and SEQ ID NO: 18) were co-transfected into COS-7 cells using a LIPOFECTIN™ based procedure as described below. COS7 cells are seeded into a 6 well plate at 2×10^5 cells/2 ml/well, from a subconfluent culture and incubated overnight at 37°, 5 % CO₂. A LIPOFECTIN™/ serum free medium mix is made up as follows: 12 ml LIPOFECTIN™ plus 200 ml serum free medium and incubated at room temperature for 30 minutes. A DNA/serum free medium mix is made up as follows: 4 mg DNA (2 mg of each construct) plus 200 ml serum free medium. 200 ml of the LIPOFECTIN™/ serum free medium mix is then added to the DNA mix and incubated for 15 minutes room temperature. 600 ml of serum free medium was then added to each sample. The cells were washed once with 2 ml serum free medium and then the 1 ml LIPOFECTIN™/DNA mix is added to the cells and incubated for 5 hours, 37°, 5 % CO₂. The LIPOFECTIN™/DNA mix was removed from the cells and normal growth media added after which the cells were incubated for 72 hours, 37°, 5 % CO₂. The cell supernatants were harvested.

g) Analysis of Antibody-Enzyme Fusion Protein

The supernatant material was analysed for the presence of antibody fusion protein using a CEA-binding ELISA using an anti human kappa light chain reporter antibody (for presence of antibody), a CEA-binding ELISA using an anti-CPG2 reporter antibody (for presence of CEA bound CPG2 fusion protein), a HPLC based CPG2 enzyme activity assay (to measure specific CPG2 activity) and SDS/PAGE followed by Western blotting (using either anti human kappa light chain reporter or anti CPG2 reporter antibodies) to detect expressed

material.

The HPLC based enzyme activity assay clearly showed CPG2 enzyme activity to be present in the cell supernatant and both the anti-CEA ELISA assays exhibited binding of protein at levels commensurate with a bivalent A5B7 antibody molecule. The fact that the anti-CEA ELISA detected with an anti-CPG2 reporter antibody also exhibited clear CEA binding indicated that not only antibody but also antibody-CPG2 fusion protein was binding CEA.

Western blot analysis with both reporter antibody assays clearly displayed a fusion protein subunit of the expected approximately 90 kDa size with no degradation or smaller products (such as Fab or enzyme) observable.

Since CPG2 is known only to exhibit enzyme activity when it is in a dimeric state and since only antibody enzyme fusion protein is present, this indicates that the 90 kDa fusion protein (seen under SDS/PAGE conditions) dimerises via the natural CPG2 dimerisation mechanism to form a 180 kDa dimeric antibody-enzyme fusion protein molecule (Figure 2a) in "native" buffer conditions. Furthermore, this molecule exhibits both CPG2 enzymatic activity and CEA antigen binding properties which do not appear to be significantly different in the fusion protein compared with enzyme or antibody alone.

h) Use of expressed fusion protein and CPG2 prodrug in an *in vitro* cytotoxicity assay

An *in vitro* cell killing assay was performed in which the (A5B7-CPG2 R6)₂ fusion protein was compared to a "conventional" A5B7 F(ab')₂-CPG2 conjugate formed through linking A5B7 F(ab')₂ to CPG2 with a chemical heterobifunctional reagent. In each case material displaying equal amounts of CPG2 enzyme activity or equal amounts of antibody-CPG2 protein were incubated with LoVo, CEA bearing, tumour cells. The cells were then washed to remove unbound protein material and subsequently resuspended in medium containing a CPG2 phenol prodrug (PGP, see Example 2 below) for a period of 1 hr, after which the cells were washed, resuspended in fresh media and left to proliferate for 4 days. Finally the cells were treated with SRB stain and their numbers determined.

The results obtained clearly showed that the (A5B7-CPG2 R6)₂ fusion protein (together with prodrug) caused at least equivalent cell kill and resulted in lower numbers of cells at the end of the assay period than the equivalent levels of A5B7 F(ab)₂-CPG2 conjugate (with the same prodrug). Cell killing (above basal control levels) can only occur if the prodrug

- 36 -

is converted to active drug by the CPG2 enzyme (and since the cells are washed to remove unbound protein, only cell bound enzyme will remain at the stage where the prodrug is added). Thus this experiment shows that at least as much of the A5B7-CPG2 R6 fusion protein remains bound compared with conventional A5B7 F(ab)₂-CPG2 conjugate as a greater degree of cell killing (presumably due to higher prodrug to drug conversion) occurs.

i) Construction of a coexpression fusion protein vector for use in transient and stable cell line expression

For a simpler transfection methodology and the direct coupling of both expression cassettes to a single selection marker, a co-expression vector for fusion protein expression was constructed using the existing vectors pNG4/A5B7VH-IgG2CH1/CPG2 R6 (encoding the antibody Fd-CPG2 fusion protein) and pNG3/A5B7VK-HuCK-NEO (encoding the antibody light chain). The pNG4/A5B7VH-IgG2CH1/CPG2 R6 plasmid was first digested with the restriction enzyme *ScaI*, the reaction loaded on a 1 % agarose gel and the linear vector band excised from the gel and purified. This vector DNA was then digested with restriction enzymes *BglII* and *BamHI*, the reaction loaded on a 1 % agarose gel, the desired band (approximately 2700 bp) recovered and purified. The plasmid pNG3/A5B7VK-HuCK-NEO was digested with the restriction enzyme *BamHI* after which the DNA was dephosphorylated then subsequently loaded on a 1 % agarose gel and the vector band excised from the gel and purified. The heavy chain expression cassette fragment was ligated in to the prepared vector and the ligation mix transformed into *E. coli*. The orientation was checked by a variety of restriction digests and clones selected which had the heavy chain cassette in the same direction as that of the light chain. These plasmids were termed pNG3-A5B7-CPG2/R6-coexp.-NEO.

j) Gene switches for protein expression

It is foreseen that *in vitro* expression of CPG2 and CPG2 fusion proteins in mammalian cells may degrade media folates leading to slow cell growth or cell death. The high activity of the CPG2 enzyme is likely to make such a folate deficiency difficult to overcome by media supplementation. However, it is thought that in the case of CPG2 or CPG2 fusion protein expression from mammalian cells *in vivo*, it is unlikely that such problems will occur, since the cells would be constantly replenished with all growth requirements by the normal circulatory and cellular mechanisms.

A number of options to avoid possible *in vitro* folic acid depletion problems have been considered. One of these solutions involve the use of tightly controlled but inducible gene switch systems such as the "TET on" or "TET off" switches (Grossen, M. et al (1995) Science 268: 1766-1769) or the ecdysone/ muristerone A switch (No, D. et al (1996) PNAS 93 :3346-3351). Such systems enable precisely controlled expression of a gene of interest and allow stable transformation of mammalian cells with genes encoding toxic or potentially deleterious expression products. A gene switch would allow recombinant stable cell lines incorporating CPG2 fusion genes to be potentially more easily established, maintained and expanded for protein expression and seeding cultures for *in vivo* tumour growth studies.

10

Example 2

HCT116 tumour cells expressing the antibody-enzyme fusion protein are selectively killed *in vitro* by a prodrug.

HCT116 colorectal tumour cells (ATCC CCL 247) transfected with the antibody-CPG2 fusion protein gene of Example 1 can be selectively killed by a prodrug that is converted by the enzyme into an active drug.

To demonstrate this, control non-transfected HCT116 cells or HCT116 cells transfected with the antibody-CPG2 fusion protein gene, are incubated with either the prodrug, 4-[N,N-bis(2-chloroethyl)amino]-phenoxycarbonyl-L-glutamic acid (PGP; Blakey et al, Br. J. cancer 72, 1083, 1995) or the corresponding drug released by CPG2, 4-[N,N-bis(2-chloroethyl)amino] phenol. PGP prodrug and drug over the concentration range of 5×10^{-4} to 5×10^{-8} M are added to 96 well microtitre plates containing 1000-2,500 HCT116 cells/well, for 1 hr at 37°. The cells are then washed and incubated for a further three days at 37°. After washing to remove dead cells, TCA is then added and the amount of cellular protein adhering to the plates is assessed by addition of SRB dye as described by Skehan et al (J. Natl. Cancer Inst. 82, 1107, 1990). Potency of the prodrug and drug is assessed by the concentration required to inhibit cell growth by 50 % (IC₅₀).

Treatment of non-transfected or transfected HCT116 cells with the drug results in an IC₅₀ of approximately 1 µM. In contrast, the PGP prodrug results in an IC₅₀ of approximately 200 µM on non-transfected cells and approximately 1 µM on transfected cells. These results demonstrate that the transfected cells which express the antibody-CPG2 fusion protein can convert the PGP prodrug into the more potent active drug while non-transfected HCT116 cells

30

are unable to convert the prodrug. Consequently the transfected HCT116 cells are over 100 fold more sensitive to the PGP prodrug in terms of cell killing compared to the non-transfected HCT116 cells. (See Example 1 j) for issues involving possible folic acid depletion in cells).

- 5 These studies demonstrate that transfecting tumour cells with a gene for an antibody-enzyme fusion protein can lead to selective tumour cell killing with a prodrug.

Example 3

Anti-tumour activity of PGP prodrug in HCT116 tumours expressing the antibody-CPG2 fusion protein.

- 10 The anti-tumour activity *in vivo* of the PGP prodrug in HCT116 tumours expressing the antibody-CPG2 fusion protein can be demonstrated as follows. HCT116 tumour cells transfected with the antibody-CPG2 fusion protein gene or control non-transfected HCT116 tumour cells are injected subcutaneously into athymic nude mice (10^7 tumour cells per
15 mouse). When the tumours are 5-7 mm in diameter the PGP prodrug is administered i.p. to the mice (3 doses at hourly intervals over 2 h in dose ranges of 5-25 mg kg⁻¹). The anti-tumour effects are judged by measuring the length of the tumours in two directions and calculating the tumour volume using the formula:

$$\text{Volume} = \pi/6 \times D^2 \times d$$

- 20 where D is the larger diameter and d is the smaller diameter of the tumour.

Tumour volume is expressed relative to the tumour volume at the time the PGP prodrug is administered. The anti-tumour activity is compared to a control group receiving either transfected or non-transfected tumour cells and PBS (170 mM NaCl, 3.4 mM KCl, 12 mM Na₂HPO₄ and 1.8 mM KH₂PO₄, pH 7.2) instead of the PGP prodrug.

- 25 Administration of PGP to HCT116 tumours established from transfected HCT116 cells results in a significant anti-tumour effect as judged by the PGP treated tumours decreasing in size compared to the PBS treated tumours and it taking a significantly longer time for the PGP treated tumours to reach 4 times their initial tumour volume compared to PBS treated tumours. In contrast, administration of PGP to HCT116 tumours established
30 from non-transfected cells resulted in no significant anti-tumour activity.

Similar studies can be used to demonstrate that the antibody-enzyme gene delivered in an appropriate vector to established HCT116 tumours produced from non-transfected

- 39 -

HCT116 cells when used in combination with the PGP prodrug can result in significant anti-tumour activity. Thus non-transfected HCT116 cells are injected into athymic nude mice (1 X 10⁷ tumour cells per mouse) and once the tumours are 5-7 mm in diameter the vector containing the antibody-enzyme fusion protein gene is injected intra-tumourally. After 1-3 5 days to allow the antibody-enzyme fusion protein to be expressed by and bind to the HCT116 tumour cells, the PGP prodrug is administered as described above. This results in significant anti-tumour activity compared to control mice receiving PBS instead of PGP prodrug.

Example 4

10 Improved Transfection of Adherent Cell lines Using supplemented FAS media and/or V-79 Feeder Cells

It was foreseen that *in vitro* expression of CPG2 and CPG2 fusion proteins in mammalian cells may degrade media folates leading to slow cell growth or cell death. FAS (folinic acid supplemented) media described herein was developed for CPG2 and CPG2 15 fusion protein expressing cell lines in order to better support the growth of such cell lines.

In preparation for transfection, adherent cell lines were cultured in normal DMEM media and passaged at least three times before transfection. V-79 (hamster lung fibroblast, obtained from MRC Radiobiology Unit, Harwell, Oxford, United Kingdom) feeder cells were cultured in normal DMEM media and passaged three times before use. For the transfection, a 20 viable count (using a haemocytometer/trypan blue staining) of the adherent cells was made and the cells plated out at 2 x 10⁵ cells per well into a 6 well plate (Costar 3516) and left for 18-24 hours for the cells to re-adhere.

For each individual transfection, 20µl of LIPOFECTIN™ was added to 80µl serum free medium and left at room temperature for 30 minutes. Plasmid DNA (2µg) of interest was 25 added to 100µl serum free medium and subsequently added to the LIPOFECTIN™ mix and left for a further 15 minutes. The individual 6 well plates were washed with 2 ml serum free medium per well to remove any serum and replaced with 800 µl of fresh serum free medium. The 200µl DNA / LIPOFECTIN™/serum free medium mixes which had been previously prepared were then added to each well of cells. The plates were incubated at 37° for 5 hours, 30 the media removed and 2 ml of fresh normal media added and incubated for a further 48 hours. The transfected cells in the 6 well plate were scraped free, the cell suspension removed and centrifuged. All the supernatant was removed and the cell pellet resuspended in 20 ml of

- 40 -

the appropriate fresh growth media (e.g. FAS DMEM media) containing the appropriate selective agent for the transfected DNA (e.g. G418). Aliquots (200 μ l) were plated per well into a 96 well plate (1.25×10^4 cells per well).

To enhance clone expansion, fibroblast feeder cells may be added to the transfected cells. Semi-confluent V-79 feeder cells were trypsinised and a viable count performed. The cells were resuspended to 1×10^6 cells/ml in a sterile glass container, irradiated using a Caesium source by exposure to 5000 rads over 12 minutes. The cells can then be stored at 4° for 24-48 hours (irradiated cells are metabolically active but will not divide, and so can act as "feeders" for other cells without contaminating the culture). The feeder cells should be plated out at 4×10^4 cells per well in a 96 well plate to produce a confluent layer for the emerging recombinant clones. Feeder cells initially adhere to the plate but with time detach and float off into the media, leaving the any recombinant clone still attached to the well. Media changes (200 μ l at time) are performed twice weekly to remove floating cells and replenish media. Colonies were allowed to develop for 10-14 days, then the supernatant screened by standard ELISA assay for fusion protein secretion.

To measure the expression rate in the case of the (A5B7-CPG2)₂ fusion gene constructs, recombinant cells were seeded out at 1×10^6 in 10 ml fresh normal culture media for exactly 24 hours. The supernatant was then removed, centrifuged to remove cell debris and assayed for fusion protein and enzyme activity by the ELISA and HPLC methods described above. The results for a number of recombinant (A5B7-CPG2)₂ fusion protein cell lines are shown below.

Cell Line	Clone	ng/ 10^6 cells/24h
HCT 116	F7	6550
	C12	3210
HCT 116	F6	15560
	C1	6151
	B3	4502
	A8	4650
	D5	630
	H9	610

- 41 -

	G11	2081
	H4	2380
	A4	1634
LoVo	B9	8370
	C1	7350
	F12	2983
	C7	10770
	G10	4140
Colo 320DM	B3	10540
	G4	4720
	B9	885
	B10	3090
	F12	35660

Example 5**Construction of a stable inducible (A5B7-CPG2)₂ fusion protein expressing tumour cell line****5 a) Construction of an inducible fusion protein expression vector**

To facilitate expression from a single inducible mammalian cell promoter, an IRES (Internal Ribosome Entry Site; see Y. Sugimoto *et al.*, Biotechnology (1994), **12**, 694-8) based version of the (A5B7-CPG2)₂ fusion protein was constructed. Construct pNG3 pNG3/A5B7VK-HuCK-NEO (A5B7 chimaeric light chain; described in Example 1b above) was used as a template for amplification of the light chain gene. The gene was amplified using oligonucleotides CME 3153 and CME 3231 (SEQ ID NOS 19 and 20). A PCR product of the expected size (approximately 700 b.p.) was purified. This product was then digested using the restriction enzymes EcoRI and BamHI and subsequently purified. The fragment was cloned into the Bluescript™ KS+ vector (prepared to receive the fragment by digestion with the same restriction enzymes, EcoRI and BamHI) after which the DNA was dephosphorylated and the larger vector band purified. The similarly restricted PCR fragment ligated in to the prepared vector and the ligation mix was transformed into *E. coli*. DNA was prepared from the clones obtained and analysed by restriction digestion to check for insertion of PCR fragment. Appropriate clones were sequenced to confirm the gene sequence. A

number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation **ASB7 Bluescript™**.

In a similar manner, the chimaeric A5B7 heavy chain was amplified by PCR from the plasmid **pNG4/ASB7VH-IgG2CH1/CPG2 R6** (described in Example 1e above) using
5 oligonucleotides CME 3151 and CME 3152 (SEQ ID NOS 21 and 22). A PCR reaction product of the expected size (approximately 1800 b.p.) was purified. This product was then digested using the restriction enzymes BamHI and Xba I after which the fragment band was purified. The fragment was also cloned into the Bluescript™ KS+ vector which had been prepared to receive the above fragment by digestion with the same restriction enzymes,
10 BamHI and XbaI, after which the DNA was dephosphorylated and the larger vector band was purified. The similarly restricted PCR fragment was ligated in to the prepared vector and the ligation mix was transformed into *E. coli*. DNA was prepared from the clones obtained and analysed by restriction digestion to check for insertion of PCR fragment. Appropriate clones were sequenced to confirm the gene sequence. A number of the clones with the correct
15 sequence were obtained and one of these clones was given the plasmid designation **Bluescript™ Fd-CPG2 R6**.

The IRES sequence was sourced from the vector **pSXLC** (described in Y. Sugimoto *et al.* Biotechnology (1994), 12, 694-8, and obtained from the authors). The IRES sequence was excised by digestion with the restriction enzymes BamHI and NcoI. A band of the
20 expected size (approximately 500 b.p.) was purified and ligated into the Bluescript™ Fd-CPG2 R6 plasmid (which had previously been prepared by restriction with the same enzymes). The ligation mix was transformed into *E. coli* and DNA was prepared from the clones obtained. The DNA was analysed by restriction digestion to check for insertion of the fragment and appropriate clones were subsequently sequenced to confirm the gene sequence.
25 A number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation **Bluescript™ IRES Fd-CPG2 R6**.

To facilitate later cloning steps, it was necessary to delete the Xba I site which had been carried over in the IRES fragment. This was performed by PCR mutagenesis with the oligonucleotide primers CME 3322 and CME 3306 (SEQ ID NOS: 23 and 24) and the
30 **Bluescript™ IRES Fd-CPG2 R6** as template DNA. A PCR reaction product of the expected size (approximately 500 b.p.) was purified, digested with the restriction enzymes BamHI and NcoI and ligated into the Bluescript™ IRES Fd-CPG2 R6 plasmid (which had previously

- 43 -

been prepared by restriction with the same restriction enzymes). The ligation mix was transformed into *E. coli* and DNA was prepared from the clones obtained. The DNA was analysed by restriction digestion to check for insertion of the fragment and appropriate clones were subsequently sequenced to confirm the gene sequence. A number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation **Bluescript™ IRES Fd-CPG2 R6-Xba del**.

The A5B7 chimaeric light chain fragment was excised from the A5B7 Bluescript™ plasmid by digestion with the restriction enzymes EcoRI and BamHI. A band of the expected size (approximately 700 b.p.) was purified, ligated into the appropriately prepared Bluescript IRES Fd-CPG2 R6-Xba del plasmid and the ligation mix was transformed into *E. coli*. DNA was prepared from the clones obtained and analysed by restriction digestion to check for insertion of the fragment. Appropriate clones were subsequently sequenced to confirm the gene sequence. A number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation **Bluescript™ A5B7 IRES Fd-CPG2 R6-Xba del**. The complete IRES based A5B7 chimaeric fusion protein gene sequence is shown in SEQ ID NO: 52.

The IRES based A5B7 chimaeric fusion protein gene was then transferred to a tetracycline regulated expression vector. Vectors for the Tet On gene expression system were obtained from Clontech. The Tetracycline switchable expression vector pTRE (otherwise known as pHUD10-3, see Gossen *et al.* (1992), PNAS, 89, 5547-51) was prepared to accept the IRES based fusion protein cassette by digestion with the restriction enzymes EcoRI and XbaI, dephosphorylated and the larger vector band purified. The IRES gene cassette was excised from the **Bluescript™ A5B7 IRES Fd-CPG2 R6-Xba del** plasmid using the same restriction enzymes. The approximately 3000 b.p. fragment obtained was ligated in to the prepared vector and the ligation mix was transformed into *E. coli*. DNA was prepared from the clones obtained and analysed by restriction digestion to check for insertion of PCR fragment. Appropriate clones were subsequently sequenced to confirm the gene sequence. A number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation **pHUD10-3/A5B7 IRES Fd-CPG2 R6**.

30 b) **Construction of a stable inducible fusion protein expressing cell line**

The standard lipofection transfection methodology (as described previously but without the use of feeder cells) was used to produce recombinant HCT116 tumour cell lines.

- 44 -

A co-transfection using 1 µg of the pHUD10-3/A5B7 IRES Fd-CPG2 R6 plasmid and 1 µg of the pTet-On transactivator expressing plasmid (from the Clontech kit) was performed and positive clones selected using FAS media containing 750 µg G418/ml.

c) **Induction studies of recombinant HCT116 inducible cell lines**

5 The clone cultures obtained were split in to duplicate 48 well plates, each containing 1×10^6 cells. The cells were grown for 48 h with one of the plates induced with 2 µg/ml doxycycline and the other acting as an non-induced control. Expression of the (A5B7-CPG2)₂ fusion protein in the cell supernatant was tested using the ELISA/ Western blot assays described in Example 1g. The results indicated that induction of fusion protein from
10 the inducible cell line by use of doxycycline could be clearly demonstrated, for example one of the clones obtained (F11), the induced cells produced 120 ng/ml of fusion protein in the supernatant whereas the non-induced cells produced only background levels of fusion protein (below 1 ng/ml).

15 **Example 6**

Cell based ELISA assay of secreted fusion protein material

Cells were seeded into 96 well plates (Becton Dickinson Biocoat™ poly-D-Lysine, 35-6461) at a density of 1×10^4 cells per well in 100 µl normal culture media and left about 40 h at 37°. 100 µl of 6 % formaldehyde was diluted in DMEM and left for 1 hour at 4°. Plates
20 were centrifuged and washed 3 times in PBS containing 0.05 % Tween™ by immersion soaking (first two washes for 2 minutes and the final wash for 5 minutes).

100 µl of doubling dilutions of cell culture supernatant containing fusion protein or chimeric A5B7 anti-CEA were added to each well as appropriate and the plates incubated overnight at 4°. The plates were washed as described above and, in the case of chimaeric
25 fusion proteins, 100 µl of 1:1000 dilution of HRP labelled anti-human kappa antibody (Sigma A-7164) was added and incubated for 2 hours at room temperature (an anti-CPG2 detection methodology can be used in the case of murine scFv fusion proteins). The plates were washed as described above and HRP detected using OPD substrate (Sigma P-8412). Colour was allowed to develop for about 5 min, stopped with 75 µl per well of 2M H₂SO₄ and OD read
30 at 490 nm.

In the case of the (A5B7-CPG2)₂ fusion protein, material was produced in the

supernatant from recombinant Colo320DM tumour cells (CEA-ve). The fusion protein content was measured by use of the CEA ELISAs described above. Increasing amounts of fusion protein were added to a number of CEA negative cell lines and the CEA positive LoVo parental line. The results shown in Figure 3 clearly show that only the CEA positive line shows increased levels of binding with increasing amounts of added fusion protein whereas the CEA negative cell lines show only constant background binding levels throughout. This clearly demonstrates that the fusion protein specifically binds and is retained on CEA positive Lovo cells.

10 Example 7

Recombinant LoVo tumour cells expressing antibody-enzyme fusion protein exhibit retention of the fusion protein on the cell surface

LoVo colorectal tumour cells transfected with the (A5B7-CPG2)₂ fusion protein gene have been shown both to secrete and to retain the fusion protein on their cell surface.

15 This can be demonstrated by comparing parental and recombinant fusion protein expressing LoVo cells under the conditions set out in the cell based ELISA assay of secreted fusion protein (Figure 4). On development of the colour reaction it could be seen that the recombinant LoVo cells had retained the expressed fusion protein (by showing a high level of colour). In control experiments, using Colo320DM fusion protein expressing cells, the assay

20 showed some retention of the expressed fusion protein (probably non-specific) and the parental LoVo cells only exhibited background activity. Positive controls in which CEA binding antibody was added to test recombinant fusion protein expressing tumour cells and to the parental LoVo controls resulted in a signal being obtained from the parental LoVo (thus demonstrating that CEA was present on the parental cells) but no increased signal from the

25 Colo320DM (CEA negative). The recombinant LoVo cells still gave such a strong initial signal that the added antibody made little difference to the overall signal obtained, which was considerably higher than any of the control experiments. Thus it appears that anti-CEA antibody enzyme-CPG2 fusion protein secreted from CEA positive tumour cell lines bind to the surface of the cells (via CEA) whereas the same protein expressed from CEA negative

30 tumours shows no such binding.

Example 8

LoVo tumour cells expressing the antibody-enzyme fusion protein are selectively killed *in vitro* by a prodrug.

LoVo colorectal tumour cells, transfected with the (A5B7-CPG2)₂ fusion protein gene,
5 can be selectively killed by a prodrug that is converted by CPG2 enzyme into an active drug.

To demonstrate this control non-transfected LoVo cells or LoVo cells transfected with
an antibody-CPG2 fusion protein gene are incubated with either the prodrug, 4-[N,N-bis(2-
chloroethyl)amino]-phenoxy carbonyl-L-glutamic acid (PGP; Blakey et al, (1995) Br. J. cancer
72, p1083) or the corresponding drug released by CPG2, 4-[N,N-bis(2-chloroethyl)amino]
10 phenol as described in Example 2 with HCT116 cells.

The transfected cells which express the antibody-CPG2 fusion protein can convert the
PGP prodrug into the more potent active drug while non-transfected LoVo cells are unable to
convert the prodrug.

These studies demonstrate that transfecting tumour cells with a gene for an antibody-
15 enzyme fusion protein can lead to selective tumour cell killing with a prodrug.

Example 9

Establishment of fusion protein expressing LoVo tumour xenografts in athymic mice

Recombinant LoVo fusion protein (A5B7-CPG2)₂ expressing tumour cells or mixes of
20 recombinant and parental LoVo cells were injected subcutaneously into athymic nude mice
(10⁷ tumour cells per mouse). The tumour growth rates for both 100 % recombinant and 20
%: 80 % mixes of recombinant:parental LoVo cells were compared to those of parental cell
only tumours. No significant differences were seen in the observed growth curves obtained
showing no corrections were required during comparisons between the cell lines. The tumour
25 growth rates observed showed that in each case for the xenograft tumours to reach a size of 10
x 10 mm takes about 12 days.

Example 10

Determination of enzyme activity in tumour xenograft samples

To act as a standard for the assay, a CPG2 enzyme standard curve was prepared in 20 % homogenate of normal tumour (parental cell tumour). Subsequent dilutions of samples
5 were made in the same 20 % homogenate of normal tumour.

Excised tumour tissue is removed from -80° storage (previously flash frozen in liquid nitrogen) and allowed to thaw. Any residual skin tissue was removed before the tumour was cut up in to small fragments with a scalpel. The tumour tissue was transferred to a preweighed tube and the weight of tumour tissue measured. PBS containing 0.2 mM ZnCl₂ solution was
10 added to each tumour sample to give a 20 % (w/v) mix, homogenised and placed on ice. Dilutions of sample tumours (in 20% normal tumour homogenate) were prepared e.g. neat, 1/10, 1/20 and 1/40.

For the standard curve, dilutions of CPG2 enzyme were made to the following concentrations to a final volume of 400 µl. Similarly, 400 µl of each of the recombinant
15 tumour sample dilutions were also prepared. After equilibration to 30°, 4 µl of 10 mM methotrexate (MTX) solution was added. The reaction was stopped after exactly 10 minutes by adding 600 µl ice cold methanol + 0.2 % TFA, centrifuged and the supernatant collected. The substrate and product in the supernatant were then separated by HPLC (using a Cation Exchange Column, HICROM™ S5SCX-100A, mobile phase = 60 % methanol, 40 % 60 mM
20 ammonium formate/ 0.1 % TFA, detection 300 nm). To calculate enzyme activity in the tumour tissue, the standard curve was plotted as units of area of methotrexate metabolite (the standards are such that only 20-30 % of the substrate is metabolised so ensuring this is not rate limiting). The test samples were analysed by comparing the unit area of metabolite against the standard curve and then multiplying by the dilution factor. Finally, making the
25 working assumption that 1 ml = 1 g the results were multiplied by 5 (as the samples were originally diluted to a 20% homogenate).

Results obtained with 20 % recombinant: 80 % parental LoVo cells expressing (A5B7 Fab-CPG2)₂ fusion protein showed the following results: tumours taken at day 5 had an average enzyme activity = 0.26 U/ g (range between 0.18-0.36 U/g) and at day 12 had an
30 average enzyme activity = 0.65 U/g (range between 0.19-1.1 U/g).

Example 11**Determination enzyme activity in plasma samples**

To act as a standard for the assay, a CPG2 enzyme standard curve was prepared in 20 % normal plasma to the following concentrations: 0.2, 0.4, 0.6, 0.8 and 1.0 U/ml. Similarly all 5 test plasma samples were also diluted to 20 % normal plasma. Further dilutions of these samples e.g. neat 1/10, 1/20 and 1/50 were also made using 20 % normal serum. 200 µl aliquots of each CPG2 standard and test sample dilutions were equilibrated to 30°. 2 µl of 10 mM MTX was added to each of the tubes and mixed well. to 30°. The reaction was stopped after exactly 10 minutes (to increase the sensitivity of the assay the incubation time can be 10 increased to 30 minutes) by adding 500 µl ice cold methanol + 0.2 % TFA and assayed for product using HPLC detection as described above in Example 10.

No activity was seen in the plasma except in the rare cases when the level of enzyme activity in the tumour exceeded 2.0 U/g, in which case the plasma enzyme levels were measured in the range of 0.013 to 0.045 U/ml.

15

Example 12**Anti-tumour activity of PGP prodrug in LoVo tumours expressing the antibody-CPG2 fusion protein.**

Recombinant LoVo (A5B7-CPG2)₂ fusion protein expressing tumour cells or mixes of 20 recombinant and parental LoVo cells were injected subcutaneously into athymic nude mice as described in Example 9.

When the tumours are 5-7 mm in diameter the PGP prodrug is administered i.p. to the mice (3 doses in DMSO/ 0.15 M sodium bicarbonate buffer at hourly intervals over 2 h in dose ranges of 40-80 mg kg⁻¹).

25 Anti-tumour effects are judged by measuring the length of the tumours in two directions and calculating the tumour volume using the formula

$$\text{Volume} = \pi/6 \times D^2 \times d$$

where D is the larger diameter and d is the smaller diameter of the tumour. Tumour volume may be expressed relative to the tumour volume at the time the PGP prodrug is administered 30 or alternatively the median tumour volumes may be calculated. The anti-tumour activity is compared to control groups receiving either transfected or non-transfected tumour cells and

buffer without PGP prodrug.

Administration of PGP to LoVo tumours established from recombinant LoVo cells or recombinant Lovo/Parental LoVo cell mixes results in a significant anti-tumour effect as shown by the PGP treated tumours decreasing in size compared with controls and it taking a significantly longer time for the PGP treated tumours to reach 4 times their initial tumour volume compared with controls (Figure 5). Administration of PGP to LoVo tumours established from non-transfected cells resulted in no significant anti-tumour activity.

Similar studies can be used to demonstrate that the antibody-enzyme gene delivered in an appropriate gene delivery vector to established LoVo tumours produced from non-transfected parental LoVo cells when used in combination with the PGP prodrug can result in significant anti-tumour activity. Thus non-transfected LoVo cells are injected into athymic nude mice (1 X 10⁷ tumour cells per mouse) and once the tumours are 5-7 mm in diameter the vector containing the antibody-enzyme fusion protein gene is injected intra-tumourally. After 1-3 days to allow the antibody-enzyme fusion protein to be expressed by, and bind to, the LoVo tumour cells, the PGP prodrug is administered as described above. This results in significant anti-tumour activity compared with controls.

Example 13

Construction of an (806.077 Fab-CPG2)₂ fusion protein

The construction of a (806.077 Fab-CPG2)₂ enzyme fusion was planned with the aim of obtaining a bivalent human carcinoembryonic antigen (CEA) binding molecule which also exhibits CPG2 enzyme activity. To this end the initial construct was designed to contain an 806.077 antibody heavy chain Fd fragment linked at its C-terminus via a flexible (G4S)₃ peptide linker to the N-terminus of the CPG2 polypeptide (as shown in Figure 1 but substituting 806.077 in place of A5B7).

The antibody 806.077 (described in International Patent Application WO 97/42329, Zeneca Limited) binds with a very high degree of specificity to human CEA. Thus the 806.077 antibody is particularly suitable for targeting colorectal carcinoma or other CEA antigen bearing cells.

In general, antibody (or antibody fragment)-enzyme conjugate or fusion proteins should be at least divalent, that is to say capable of binding at least 2 tumour associated antigens (which may be the same or different). In the case of the (806.077 Fab-CPG2)₂ fusion

- 50 -

protein, dimerisation of the enzyme component takes place (after expression, as with the native enzyme) thus forming an enzymatic molecule which contains two Fab antibody fragments (and is thus bivalent with respect to antibody binding sites) and two molecules of CPG2 (Figure 2a).

5 a) Cloning of the 806.077 antibody genes

Methods for the cloning and characterisation of recombinant murine 806.077 F(ab')₂ antibody have been published (International Patent Application WO 97/42329, Example 7). Reference Example 7.5, describes cloning of the 806.077 antibody variable region genes into Bluescript™ KS+ vectors. These vectors were subsequently used as the source of the 806.077
10 variable region genes for the construction of 806.077 chimaeric light and heavy chain Fd genes.

b) Chimaeric 806.077 antibody vector constructs

International Patent Application WO 97/42329, Example 8 describes the cloning of the 806.077 chimaeric light and heavy chain Fd genes in the vectors pNG3-Vkss-HuCK-NEO
15 (NCIMB deposit no. 40799) and pNG4-VHss-HuIgG2CH1' (NCIMB deposit no. 40797) respectively. The resulting vectors were designated pNG4/VHss806.077VH-IgG2CH1' (806.077 chimaeric heavy chain Fd') and pNG3/VKss806.077VK-HuCK-NEO (806.077 chimaeric light chain). These vectors were the source of the 806.077 antibody genes for the construction of the 806.077 Fab-CPG2 fusion protein.

20 c) Construction of the 806.077 heavy chain Fd-CPG2 fusion protein gene

The cloning and construction of the CPG2 gene used are described in Example 1, sections c and d. Similarly, the construction of the pNG4/A5B7VH-IgG2CH1/CPG2 R6 vector, which was used for the construction of the 806.077 heavy chain Fd-CPG2, is described in Example 1, section e. The 806.077 variable heavy chain gene was removed from the
25 pNG4/VHss806.077VH-IgG2CH1' vector by digestion with restriction enzymes HindIII and NheI and a band of the expected size (approximately 300 b.p) which contained the variable region gene was purified. The same restriction enzymes (HindIII/NheI) were used to digest the vector pNG4/A5B7VH-IgG2CH1/CPG2 R6 in preparation for the substitution of the 806.077 variable region for that of the A5B7 antibody. After digestion, the DNA was
30 dephosphorylated then the larger vector band was separated and purified. The similarly restricted variable region gene fragment was then ligated in to this prepared vector and the ligation mix transformed into *E. coli*. DNA was prepared from the clones obtained and

analysed by restriction digest analysis and subsequently sequenced to confirm the fusion gene sequence. A number of the clones were found to be correct and one of these clones, pNG4/VHss806VH-IgG2CH1/CPG2 R6, was chosen for further work. The sequence of the 806.077 heavy chain Fd-CPG2 fusion protein gene created is shown SEQ ID NOS 25 and 26.

5 d) **Co-transfection, transient expression and analysis of fusion protein**

The plasmids pNG4/VHss806.077VH-IgG2CH1/CPG2 R6 (encoding the antibody chimaeric Fd-CPG2 fusion protein) and pNG3/VHss806.077VK-HuCK-NEO (encoding the antibody chimaeric light chain) were co-transfected into COS-7 cells using a LIPOFECTIN™ based procedure described in Example 1f above. Analysis of the fusion protein was
10 performed as described in Example 1g. The HPLC based enzyme activity assay clearly showed CPG2 enzyme activity to be present in the cell supernatant and both the anti-CEA ELISA assays exhibited binding of protein at levels commensurate with a bivalent 806.077 antibody molecule. The fact that the anti-CEA ELISA detected with an anti-CPG2 reporter antibody also exhibited clear CEA binding indicated that not only antibody but also antibody-
15 CPG2 fusion protein was binding CEA. Western blot analysis with both reporter antibody assays clearly displayed a (806.077 Fab-CPG2)₂ fusion protein subunit of the expected approximately 90 kDa size with only a small amount of degradation or smaller products (such as Fab or enzyme) observable. Since CPG2 is only known to exhibit enzyme activity when it is in a dimeric state it and since only antibody enzyme fusion protein is present, this indicates
20 that the 90 kDa fusion protein (seen under SDS/PAGE conditions) dimerises via the natural CPG2 dimerisation mechanism to form a 180 kDa dimeric antibody-enzyme fusion protein molecule (Figure 2a) in "native" buffer conditions. Furthermore, this molecule exhibits both CPG2 enzymatic activity and CEA antigen binding properties which do not appear to be significantly different in the fusion protein compared with enzyme or antibody alone.

25 e) **Construction of a (806.077 Fab-CPG2)₂ fusion protein coexpression vector for use in transient and stable cell line expression**

For a simpler transfection methodology and the direct coupling of both expression cassettes to a single selection marker, a co-expression vector for fusion protein expression was constructed using the existing vectors pNG4/VHss806.077VH-IgG2CH1/CPG2 (encoding
30 the antibody Fd-CPG2 fusion protein) and pNG3/VKss806.077VK-HuCK-NEO (encoding the antibody light chain). The pNG4/VHss806.077VH-IgG2CH1/CPG2 plasmid was first digested with the restriction enzyme ScaI, the linear vector band purified, digested with the

- 52 -

restriction enzymes BglII and BamHI and a desired band (approximately 2700 b.p.) purified. The plasmid pNG3/VKss806.077VK-HuCK-NEO was digested with the restriction enzyme BamHI after which the DNA was dephosphorylated and the vector band purified. The heavy chain expression cassette fragment was ligated in to the prepared vector and the ligation mix transformed into *E. coli*. The orientation was checked by a variety of restriction digests and clones selected which had the heavy chain cassette in the same direction as that of the light chain. This plasmid was termed **pNG3-806.077-CPG2/R6-coexp.-NEO**.

Example 14

10 Construction of a (55.1 scFv-CPG2)₂ fusion protein

The 55.1 antibody, described in the United States Patent 5,665,357, recognises the CA55.1 tumour associated antigen which is expressed on the majority of colorectal tumours and is only weakly expressed or absent in normal colonic tissue. The determination of the 55.1 heavy and light chain cDNA sequences is described in Example 3 of the aforementioned US patent. A plasmid expression vector allowing the secretion of antibody fragments into the periplasm of *E. coli* utilizing a single pelB leader sequence (**pICI266**) has been deposited as accession number NCIMB 40589 on 11 Oct 93 under the Budapest Treaty at the National Collections of Industrial and Marine Bacteria Limited (NCIMB), 23 St. Machar Drive, Aberdeen, AB2 1RY, Scotland, U.K. This vector was modified as described in Example 3.3a of United States Patent 5,665,357 to create **pICI1646**; this plasmid was used for cloning of various 55.1 antibody fragments as described in further subsections of Example 3, including the production of a 55.1 scFv construct which was designated **pICI1657**.

The **pICI1657** (otherwise known as **pICI-55.1 scFv**) was used as the starting point for the construction of the (55.1 scFv-CPG2)₂ fusion protein. The 55.1 scFv gene was amplified using the oligonucleotides CME 3270 and CME 3272 (SEQ ID NOS: 27 and 28 respectively) and the plasmid **pICI1657** as the template DNA. The resulting PCR product band of about 790 b.p. was purified. Similarly the pNG4/A5B7VH-IgG2CH1/CPG2 R6 plasmid described in Example 1e above was used as the template DNA in a standard PCR reaction to amplify the CPG2 gene using the oligonucleotide primers CME 3274 and CME 3275 (SEQ ID NOS: 29 and 30 respectively). The expected PCR product band of about 1200 b.p. was purified.

A further PCR reaction was performed to join (or splice) the two purified PCR reaction products together. Standard PCR reaction conditions were used using varying

- 53 -

amounts (between 0.5 to 2 μ l) of each PCR product but utilising 25 cycles (instead of the usual 15 cycles) with the oligonucleotides CME 3270 and CME 3275 (SEQ ID NOS: 27 & 30). A reaction product of the expected size (approximately 2000 b.p.) was excised, purified and eluted in 20 μ l H₂O, digested using the restriction enzyme EcoRI and purified. The

5. vector pNG4/VHss806.077VH-IgG2CH1/CPG2 was prepared to receive the above PCR product by digestion with restriction enzyme EcoRI, dephosphorylated, the larger vector band separated from the smaller fragment and purified. The similarly restricted PCR product was ligated in to the prepared vector and the ligation mix transformed into *E. coli*. DNA was prepared from the clones obtained and analysed by HindIII/NotI restriction digestion to check

10 for correct fragment orientation and appropriate clones subsequently sequenced to confirm the fusion gene sequence. A number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation pNG4/55.1scFv/CPG2 R6. The DNA and amino acid sequences of the fusion protein are shown in SEQ ID NOS: 31 and 32.

15 Example 15

Modification of the plasmid pNG4/55.1scFv/CPG2 R6 to facilitate scFv gene exchange

During the construction of pNG4/55.1scFv/CPG2 R6 a unique BspEI (isoschizomer of AccIII) was introduced into the flexible (G₄S)₃ linker coding sequence, situated between the antibody and CPG2 genes. To facilitate cloning of alternative scFv constructs the EcoRI

20 site 3' of the CPG2 gene in the pNG4/55.1scFv/CPG2 R6 was deleted in order to enable insertion of alternative scFv antibody genes in frame, both behind the plasmid signal sequence and 5' of the CPG2 gene, via a EcoRI/BspEI fragment cloning. This modification was achieved by PCR mutagenesis in which first the pNG4/55.1scFv/CPG2 R6 was amplified using oligonucleotides CME 3903 and CME 3906 (SEQ ID NOS: 33 and 34 respectively).

25 Secondly, the pNG4/55.1scFv/CPG2 R6 was again amplified but using oligonucleotides CME 4040 and CME 3905 (SEQ ID NOS: 35 and 36 respectively). The first expected PCR product band of about 420 b.p. was purified. The second PCR reaction was similarly treated and the expected PCR product band of about 450 b.p. purified.

A further PCR reaction was performed to join (or splice) the two purified PCR

30 reaction products together. Standard PCR reaction conditions were used using varying amounts (between 0.5 to 2 μ l) of each PCR product but utilising between 15 and 25 cycles

- 54 -

with oligonucleotides CME 3905 and CME 3906 (SEQ ID NOS: 36 & 34). A reaction product of the expected size (approximately 840 b.p.) was purified, digested using the restriction enzymes NotI and XbaI and the expected fragment band of ca.460 b.p. was purified.

- 5 The original pNG4/55.1scFv/CPG2 R6 was prepared to receive the above PCR product by digestion with restriction enzymes NotI and XbaI, dephosphorylated and the larger vector band separated from the smaller fragment. The vector band was purified and subsequently the similarly restricted PCR product was ligated in to the prepared vector and the ligation mix transformed into *E. coli*. DNA was prepared from the clones obtained and
- 10 analysed by EcoRI restriction digestion to check for insertion of the modified fragment and appropriate clones subsequently sequenced to confirm the sequence change. A number of clones with the correct sequence were obtained and one of these clones was given the plasmid designation pNG4/55.1scFv/CPG2 R6/del EcoRI. This mutation removes the EcoRI site which was 3' of the CPG2 gene and simultaneously introduces an additional stop codon. The
- 15 DNA sequence of the fusion protein gene up to, and including the two stop codons, are shown in SEQ ID NO: 37.

Example 16

Construction of an 806.077 scFv antibody gene

- 20 The 806.077 scFv was created using vectors pNG4/VHss806.077VH-IgG2CH1' and pNG3/VKss806.077VK-HuCK-NEO which are sources for 806.077 VH and VK variable region genes. The 806.077 VH gene was amplified from the pNG4/VHss806.077VH-IgG2CH1' plasmid using standard PCR conditions with the oligonucleotides CME 3260 and CME 3266 (SEQ ID NOS: 39 and 40 respectively). The 806.077 VK was amplified from the
- 25 pNG3/VKss806.077VK-HuCK-NEO plasmid using oligonucleotides CME 3262 and CME 3267 (SEQ ID NOS: 41 and 42 respectively). The VH and VK PCR reaction products were purified.

- A further PCR reaction was performed to join (or splice) the two purified PCR reaction products together. Standard PCR reaction conditions were used using varying
- 30 amounts (between 0.5 to 2 µl) of each PCR product but utilising between 15 and 25 cycles with the flanking oligonucleotides oligonucleotides CME 3260 and CME 3262 (SEQ ID NOS: 39 & 41). A reaction product of the expected size (approximately 730 b.p.) was

- 55 -

purified, digested using the restriction enzymes NcoI and XhoI and an expected fragment band of about 720 b.p. purified.

The pICI1657 plasmid (otherwise known as pICI-55.1 scFv) had been further modified by the insertion of a double stranded DNA cassette produced from the two oligonucleotides CME 3143 and CME 3145 (SEQ ID NOS: 45 and 46) between the existing XhoI and EcoR restriction sites by standard cloning techniques to create the vector **pICI266-55.1 scFv tag/his** (the DNA sequence of the resulting 55.1 scFv tag/his gene is shown in SEQ ID NO: 47). This vector was prepared to receive the above PCR product by digestion with restriction enzymes NcoI and XhoI, dephosphorylated and the larger vector band separated from the smaller fragment. The vector band was purified and subsequently the similarly restricted PCR product was ligated in to the prepared vector and the ligation mix transformed into *E. coli*. DNA was prepared from the clones obtained and analysed by EcoRI restriction digestion to check for insertion of the modified fragment and appropriate clones subsequently sequenced to confirm the sequence change. A number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation **pICI266/806IscFvtag/his** (alternatively known as pICI266-806VH/VLscFvtag/his). The DNA and protein sequences of the 806I scFvtag/his gene are shown in (SEQ ID NOS: 25 and 26).

20 Example 17

Construction of an (806.077 scFv-CPG2)₂ fusion protein

The pICI266/806IscFvtag/his plasmid was used as the source for the 806scFv. The gene was amplified using oligonucleotides CME 3907 and CME 3908 (SEQ ID NOS: 48 and 49) and a band of the expected size purified. This fragment was then digested using the restriction enzymes EcoRI and BspEI after which an expected fragment band of about 760 b.p. was purified.

The pNG4/55.1scFv/CPG2 R6/del EcoRI plasmid was prepared to receive the above fragment by digestion with restriction enzymes EcoRI and BspEI, dephosphorylated and the larger vector band separated from the smaller fragment. The vector band was purified and subsequently the similarly restricted fragment ligated in to the prepared vector and the ligation mix was transformed into *E. coli*. DNA was prepared from the clones obtained and analysed by EcoRI restriction digestion to check for insertion of the modified fragment. Appropriate

- 56 -

clones were subsequently sequenced to confirm the gene sequence. A number of the clones with the correct sequence were obtained and one of these clones was given the plasmid designation pNG4/806IscFv/CPG2 R6/del EcoRI. The DNA and protein sequence of the fusion protein gene 806IscFv/CPG2 R6 are shown in (SEQ ID NOS: 50 and 51).

5

Example 18

Co-transfection, transient expression of antibody-CPG2 fusion proteins

As described in Example 1f, plasmids encoding other fusion protein variants can be transfected using the given standard conditions in order to obtain transient expression of their encoded fusion protein from COS7 cells. In the case of (Fab-CPG2)₂ fusion proteins both co-transfection of appropriate plasmids or transfection of co-expression proteins can be performed. Similarly, the single expression plasmids of (scFv-CPG2)₂ fusion proteins can be also be transfected by the same protocol. In each case a maximum total of 4 mg DNA are used in an individual transfection.

15

Example 19

Gene switches for protein expression

As described in Example 1j, the use of tightly controlled but inducible gene switch systems such as the "TET on" or "TET off" (Grossen, M. et al (1995) Science 268: 1766-1769) or the ecdysone/ muristerone A (No, D. et al (1996) PNAS 93 :3346-3351) may be used for the expression of fusion proteins. Appropriate methodology and cloning strategies as described in Example 5 may be used for antibody Fab-enzyme fusions requiring an IRES sequence for expression. Insertion of the appropriate gene cassette in to the switchable expression vectors may be used if the fusion protein product is a single polypeptide chain such as in scFv-enzyme constructs.

25

Example 20

Determination of the properties of COS7 cell secreted antibody-enzyme fusion proteins

The COS7 cell supernatant material can be analysed for the presence of antibody fusion proteins as described in Example 1g. Similarly the use of expressed fusion protein and CPG2 prodrug in an *in vitro* cytotoxicity assay can be performed as previously described in Example 1h. The HPLC based enzyme activity assay can show CPG2 enzyme activity to be

30

- 57 -

present in the cell supernatant and anti-CEA ELISA can be detected with an anti-CPG2 reporter antibody to confirm binding of protein at levels commensurate with a bivalent A5B7 antibody molecule and also to demonstrate that antibody-CPG2 fusion protein (not only just the antibody component) is binding CEA.

5 Western blot analysis with both reporter antibody assays clearly display a fusion protein subunit of the expected size. Since CPG2 is only known to exhibit enzyme activity when it is in a dimeric state it and since only antibody enzyme fusion protein is present, this indicates that the fusion protein (seen under SDS/PAGE conditions) dimerises via the natural CPG2 dimerisation mechanism to form a dimeric antibody-enzyme fusion protein molecule in
10 "native" buffer conditions. Furthermore, this molecule exhibits both CPG2 enzymatic activity and CEA antigen binding properties which do not appear to be significantly different in the fusion protein compared with enzyme or antibody alone. Results obtained from the cytotoxicity assay can demonstrate that antibody-enzyme fusion protein (together with prodrug) causes at least equivalent cell kill and resulted in lower numbers of cells at the end of
15 the assay period than the equivalent levels of A5B7 F(ab')₂-CPG2 conjugate (with the same prodrug). Since cell killing (above basal control levels) can only occur if the prodrug is converted to active drug by the CPG2 enzyme (and since the cells are washed to remove unbound protein, only cell bound enzyme will remain at the stage where the prodrug is added). Thus this experiment can demonstrate that at least as much of the (A5B7-CPG2 R6)₂
20 fusion protein remains bound compared with conventional A5B7 F(ab)₂-CPG2 conjugate as a greater degree of cell killing (presumably due to higher prodrug to drug conversion) occurs.

Example 21

***In vitro* and *in vivo* determination of the properties of antibody-enzyme fusion proteins 25 expressed from recombinant tumour cells**

The construction of fusion protein expressing tumour cell lines can be performed as described in Example 4.

Retention of the fusion protein on the cell surface of recombinant LoVo tumour cells expressing antibody-enzyme fusion protein can be shown using the techniques described in
30 Example 7. Selective killing of cultured LoVo tumour cells transfected with an antibody-CPG2 fusion protein gene by a prodrug that is converted by the enzyme into an active drug

can be demonstrated as described in Example 8. Establishment of antibody-enzyme fusion protein expressing LoVo tumours xenografts in athymic mice can be performed as described in Example 9. Determination of enzyme activity in tumour xenograft samples can also be determined as described in Example 10.

- 5 Determination enzyme activity in plasma samples performed as described in Example 11. The anti-tumour activity of PGP prodrug in LoVo tumours expressing the antibody-CPG2 fusion protein can be evaluated using the method described in Example 12.

 The results from these experiments can be used to show that the antibody-CPG2 fusion protein secreted from CEA positive tumour cell lines bind to the surface of the cells
10 (via CEA) whereas the same protein expressed from CEA negative tumours shows no such binding.

 These results can demonstrate that the transfected cells which express the antibody-CPG2 fusion protein can convert the PGP prodrug into the more potent active drug while non-transfected LoVo cells are unable to convert the prodrug. Consequently the transfected LoVo
15 cells will be over 100 fold more sensitive to the PGP prodrug in terms of cell killing compared to the non-transfected LoVo cells, thus demonstrating that transfecting tumour cells with a gene for an antibody-enzyme fusion protein can lead to selective tumour cell killing with a prodrug.

 Administration of PGP to LoVo tumours established from recombinant LoVo cells or
20 recombinant Lovo/Parental LoVo cell mixes can result in a significant anti-tumour effect as judged by the PGP treated tumours decreasing in size compared to the formulation buffer only treated tumours and it taking a significantly longer time for the PGP treated tumours to reach 4 times their initial tumour volume compared with formulation buffer treated tumours. In contrast, administration of PGP to LoVo tumours established from non-transfected cells
25 would result in no significant anti-tumour activity.

 Similar studies can be used to demonstrate that the antibody-enzyme gene delivered in an appropriate gene delivery vector to established LoVo tumours produced from non-transfected parental LoVo cells when used in combination with the PGP prodrug can result in significant anti-tumour activity. Thus non-transfected LoVo cells are injected into athymic
30 nude mice (1×10^7 tumour cells per mouse) and once the tumours are 5-7 mm in diameter the vector containing the antibody-enzyme fusion protein gene is injected intra-tumourally. After 1-7 days to allow the antibody-enzyme fusion protein to be expressed by, and bind to, the

LoVo tumour cells, the PGP prodrug is administered as previously described. This results in significant anti-tumour activity compared with control mice receiving formulation buffer instead of PGP prodrug.

5 Example 22

Preparation of (murine A5B7 Fab-CPG2)₂ fusion protein

(Murine A5B7 Fab-CPG2)₂ is expressed from COS-7 and CHO cells essentially as described in part (d) of Example 48 of International Patent Application WO 97/42329 (Zeneca Limited, published 13 November, 1997) by cloning the genes for A5B7 light chain and A5B7
10 Fd linked at its C-terminus via a flexible (G₄S)₃ peptide linker to CPG2 in the pEE14 co-expression vector.

The murine A5B7 light chain is isolated from pAF8 (described in part g of Reference Example 5 in International Patent Application WO 96/20011, Zeneca Limited). Plasmid pAF8 is cut with EcoRI and the resulting 732 bp fragment isolated by electrophoresis on a 1%
15 agarose gel. This fragment is cloned into pEE14 (described by Bebbington in METHODS: A Companion to Methods in Enzymology (1991) 2, 136-145) similarly cut with EcoRI and the resulting plasmid used to transform *E. coli* strain DH5 α . The transformed cells are plated onto L agar plus ampicillin (100 μ g/ml). A clone containing a plasmid with the correct sequence and orientation is confirmed by DNA sequence analysis (SEQ ID NO: 57) and the
20 plasmid named pEE14/A5B7muVkmucK. The amino acid sequence of the encoded signal sequence (amino acid residues 1 to 22) and murine light chain (amino acid residues 23 to 235) is shown in SEQ ID NO: 58.

The murine Fd-CPG2 gene is prepared from the R6 variant of the CPG2 gene (d of Example 1) and the murine A5B7 Fd sequence in pAF1 (described in part d of Reference
25 Example 5 in International Patent Application WO 96/20011, Zeneca Limited). A PCR reaction with oligonucleotides SEQ ID NOS: 53 and 54 on pAF1 gives a 247 bp fragment. This is cut with HindIII and BamHI and cloned into similarly cut pUC19. The resulting plasmid is used to transform *E. coli* strain DH5 α . The transformed cells are plated onto L agar plus ampicillin (100 μ g/ml). A clone containing a plasmid with the correct sequence is
30 named pUC19/muCH1/NcoI-AccIII(Fd). A second PCR with oligonucleotides SEQ ID NOS: 55 and 56 on pNG/VKss/CPG2/R6-neo (Example 1) gives a 265 bp fragment which is cut with HindIII and EcoRI and cloned into similarly cut pUC19 as above to give plasmid

- 60 -

pUC19/muCH1-linker-CPG2/AccIII-SacII. Plasmid pUC19/muCH1/NcoI-AccIII(Fd) is cut with HindIII and AccIII and the 258 bp fragment isolated by electrophoresis on a 1 % agarose gel. This fragment is cloned into HindIII and AccIII cut pUC19/muCH1-linker-CPG2/AccIII-SacII to give plasmid pUC19/muCH1-linker-CPG2/NcoI-SacII. A 956 bp fragment is
 5 isolated from pNG/VKss/CPG2/R6-neo by cutting it with SacII and EcoRI. This is cloned into SacII and EcoRI cut pUC19/muCH1-linker-CPG2/NcoI-SacII to give plasmid pUC19/muCH1-linker-RC/CPG2(R6). The complete gene construct is prepared by isolating a 498 bp HindIII to NcoI fragment from pAF1 and cloning it into HindIII and NcoI cut pUC19/muCH1-linker-RC/CPG2(R6). The resulting plasmid is used to transform *E. coli*
 10 strain DH5 α . The transformed cells are plated onto L agar plus ampicillin (100 μ g/ml). A clone containing a plasmid with the correct sequence and orientation is confirmed by DNA sequence analysis (SEQ ID NO: 59) and the plasmid named pUC19/muA5B7-RC/CPG2(R6). The amino acid sequence of the encoded signal sequence (amino acid residues 1 to 19) and murine Fd-linker-CPG2 (amino acid residues 20 to 647) is shown in SEQ ID NO: 60.
 15 Alternatively, the CPG2 gene sequence described in Example 1 can be obtained by total gene synthesis and converted to the R6 variant as described in d of Example 1. In this case, the base residue C at position 933 in SEQ ID NO: 59 is changed to G. The amino acid sequence of SEQ ID NO: 60 remains unaltered.

For expression in the pEE14 vector, the gene is first cloned into pEE6 (this is a
 20 derivative of pEE6.hCMV - Stephens and Cockett, 1989, Nucleic Acids Research 17, 7110, in which a HindIII site upstream of the hCMV promoter has been converted to a BglII site). Plasmid pUC19/muA5B7-RC/CPG2(R6) is cut with HindIII and EcoRI and the 1974 bp fragment isolated by electrophoresis on a 1 % agarose gel. This is cloned into HindIII and EcoRI cut pEE6 in *E. coli* strain DH5 α to give plasmid pEE6/muA5B7-RC/CPG2(R6). The
 25 pEE14 co-expression vector is made by first cutting pEE6/muA5B7-RC/CPG2(R6) with BglII and BamHI and isolating the 4320 bp fragment on a 1 % agarose gel. This fragment is cloned into BglII and BamHI cut pEE14/A5B7muVkmucK. The resulting plasmid is used to transform *E. coli* strain DH5 α . The transformed cells are plated onto L agar plus ampicillin (100 μ g/ml). A clone containing a plasmid with the correct sequence and orientation is
 30 confirmed by DNA sequence analysis and the plasmid named pEE14/muA5B7-RC/CPG2(R6).

- 61 -

For expression of (murine A5B7 Fab-CPG2)₂, plasmid pEE14/muA5B7-RC/CPG2(R6) is used to transfect COS-7 or CHO cells as described in Example 48 of International Patent Application WO 97/42329, Zeneca Limited, published 13 November 1997. COS cell supernatants and CHO clone supernatants are assayed for activity as described in Example 1 and shown to have CEA binding and CPG2 enzyme activity.

Example 23

Pharmaceutical composition

The following illustrate a representative pharmaceutical dosage form containing a gene construct of the invention which may be used for therapy in combination with a suitable prodrug.

A sterile aqueous solution, for injection either parenterally or directly into tumour tissue, containing 10^7 - 10^{11} adenovirus particles comprising a gene construct as described in Example 1. After 3-7 days, three 1 g doses of prodrug are administered as sterile solutions at hourly intervals. Prodrug is selected from N-(4-[N,N-bis(2-iodoethyl)amino]-phenoxy-carbonyl)-L-glutamic acid, N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic- γ -(3,5-dicarboxy)anilide or N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic acid or a pharmaceutically acceptable salt thereof.

- 62 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT:

- (A) NAME: Zeneca Limited
- (B) STREET: 15 Stanhope Gate
- (C) CITY: London
- 10 (D) STATE: England
- (E) COUNTRY: United Kingdom
- (F) POSTAL CODE (ZIP): W1Y 6LN
- (G) TELEPHONE: 0171 304 5000
- (H) TELEFAX: 0171 304 5151
- 15 (I) TELEX: 0171 304 2042

(ii) TITLE OF INVENTION: CHEMICAL COMPOUNDS

20

(iii) NUMBER OF SEQUENCES: 60

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- 25 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: GB 9709421.3
- (B) FILING DATE: 10-MAY-1997

30

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GGGAATTCCT CGAGGAGCTC C

21

45 (2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs

- 63 -

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CCGGGGAGCT CCTCGAGGAA TTCCCGC

27

- 10 (2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

- 15 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CAGAAGCGCG ACAACGTG

18

20

- (2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

25

- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

30 CGAGGCCTTG CCGGTGATCT GGACCTGCAC GTAGGCGAT

39

- (2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

40

GGGGATGATG TTCGAGACCT GGCCGGCCTT GGCGATGGTC CACTGGAAGC GCAGGTTCTT
CGC

60

63

- (2) INFORMATION FOR SEQ ID NO: 6:

45

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid

- 64 -

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: other nucleic acid
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:
- 5 CTTGCCGGCG CCCAGATC 18
- (2) INFORMATION FOR SEQ ID NO: 7:
- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: other nucleic acid
15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
- GTCTCGAACA TCATCCCC 18
- (2) INFORMATION FOR SEQ ID NO: 8:
- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
25 (ii) MOLECULE TYPE: other nucleic acid
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:
- ATCACCGGCA AGGCCTCG 18
- 30 (2) INFORMATION FOR SEQ ID NO: 9:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1236 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
35 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: other nucleic acid
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
- ATGGATTTTC AAGTGCAGAT TTTCAGCTTC CTGCTAATCA GTGCTTCAGT CATAATGTCC 60
40 CGCGGGCAGA AGCGCGACAA CGTGCTGTTC CAGGCAGCTA CCGACGAGCA GCCGGCCGTG 120
ATCAAGACGC TGGAGAAGCT GGTCAACATC GAGACCGGCA CCGGTGACGC CGAGGGCATC 180
GCCGCTGCGG GCAACTTCCT CGAGGCCGAG CTCAGAACC TCGGCTTCAC GGTACGCGA 240
AGCAAGTCGG CCGGCCTGGT .GGTGGCGAC AACATCGTGG GCAAGATCAA GGGCCGCGG 300
GGCAAGAACC TGCTGCTGAT GTCGCACATG GACACCGTCT ACCTCAAGGG CATTCTCGCG 360
45 AAGGCCCCGT TCCGCGTCGA AGGCGACAAG GCCTACGGCC CGGGCATCGC CGACGACAAG 420
GGCGGCAACG CGGTCATCCT GCACACGCTC AAGCTGCTGA AGGAATACGG CGTGCGCGAC 480
TACGGCACCA TCACCGTGCT GTTCAACACC GACGAGGAAA AGGGTTCCTT CGGCTCGCGC 540

- 65 -

GACCTGATCC AGGAAGAAGC CAAGCTGGCC GACTACGTGC TCTCCTTCGA GCCCACCAGC 600
 GCAGGCGACG AAAAAGTCTC GCTGGGCACC TCGGGCATCG CCTACGTGCA GGTCCAGATC 660
 ACCGGCAAGG CCTCGCATGC CGGCGCCGCG CCCGAGCTGG GCGTGAACGC GCTGGTCGAG 720
 GCTTCCGACC TCGTGCTGCG CACGATGAAC ATCGACGACA AGGCGAAGAA CCTGCGCTTC 780
 5 CAGTGGACCA TCGCCAAGGC CGGCCAGGTC TCGAACATCA TCCCCGCCAG CGCCACGCTG 840
 AACGCCGACG TCGCTACGC GCGCAACGAG GACTTCGACG CCGCCATGAA GACGCTGGAA 900
 GAGCGCGCGC AGCAGAAGAA GCTGCCCAG GCGACGTGA AGGTGATCGT CACGCGCGGC 960
 CGCCCGGCCT TCAATGCCCG CGAAGGCGGC AAGAAGCTGG TCGACAAGGC GGTGGCCTAC 1020
 TACAAGGAAG CCGGCGGCAC GCTGGGCGTG GAAGAGCGCA CCGGCGGCGG CACCGACGCG 1080
 10 GCCTACGCCG CGCTCTCAGG CAAGCCAGTG ATCGAGAGCC TGGGCGTGCC GGGCTTCGGC 1140
 TACCACAGCG ACAAGGCCGA GTACGTGGAC ATCAGCGCGA TTCCGCGCCG CCTGTACATG 1200
 GCTGCGCGCC TGATCATGGA TCTGGGCGCC GGCAAG 1236

(2) INFORMATION FOR SEQ ID NO: 10:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 412 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

20

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

25 Met Asp Phe Gln Val Gln Ile Phe Ser Phe Leu Leu Ile Ser Ala Ser
 1 5 10 15
 Val Ile Met Ser Arg Gly Gln Lys Arg Asp Asn Val Leu Phe Gln Ala
 20 25 30
 Ala Thr Asp Glu Gln Pro Ala Val Ile Lys Thr Leu Glu Lys Leu Val
 35 40 45
 30 Asn Ile Glu Thr Gly Thr Gly Asp Ala Glu Gly Ile Ala Ala Ala Gly
 50 55 60
 Asn Phe Leu Glu Ala Glu Leu Lys Asn Leu Gly Phe Thr Val Thr Arg
 65 70 75 80
 Ser Lys Ser Ala Gly Leu Val Val Gly Asp Asn Ile Val Gly Lys Ile
 35 85 90 95
 Lys Gly Arg Gly Gly Lys Asn Leu Leu Leu Met Ser His Met Asp Thr
 100 105 110
 Val Tyr Leu Lys Gly Ile Leu Ala Lys Ala Pro Phe Arg Val Glu Gly
 115 120 125
 40 Asp Lys Ala Tyr Gly Pro Gly Ile Ala Asp Asp Lys Gly Gly Asn Ala
 130 135 140
 Val Ile Leu His Thr Leu Lys Leu Leu Lys Glu Tyr Gly Val Arg Asp
 145 150 155 160
 Tyr Gly Thr Ile Thr Val Leu Phe Asn Thr Asp Glu Glu Lys Gly Ser
 45 165 170 175
 Phe Gly Ser Arg Asp Leu Ile Gln Glu Glu Ala Lys Leu Ala Asp Tyr
 180 185 190

- 66 -

Val Leu Ser Phe Glu Pro Thr Ser Ala Gly Asp Glu Lys Leu Ser Leu
 195 200 205
 Gly Thr Ser Gly Ile Ala Tyr Val Gln Val Gln Ile Thr Gly Lys Ala
 210 215 220
 5 Ser His Ala Gly Ala Ala Pro Glu Leu Gly Val Asn Ala Leu Val Glu
 225 230 235 240
 Ala Ser Asp Leu Val Leu Arg Thr Met Asn Ile Asp Asp Lys Ala Lys
 245 250 255
 Asn Leu Arg Phe Gln Trp Thr Ile Ala Lys Ala Gly Gln Val Ser Asn
 10 260 265 270
 Ile Ile Pro Ala Ser Ala Thr Leu Asn Ala Asp Val Arg Tyr Ala Arg
 275 280 285
 Asn Glu Asp Phe Asp Ala Ala Met Lys Thr Leu Glu Glu Arg Ala Gln
 290 295 300
 15 Gln Lys Lys Leu Pro Glu Ala Asp Val Lys Val Ile Val Thr Arg Gly
 305 310 315 320
 Arg Pro Ala Phe Asn Ala Gly Glu Gly Gly Lys Lys Leu Val Asp Lys
 325 330 335
 Ala Val Ala Tyr Tyr Lys Glu Ala Gly Gly Thr Leu Gly Val Glu Glu
 20 340 345 350
 Arg Thr Gly Gly Gly Thr Asp Ala Ala Tyr Ala Ala Leu Ser Gly Lys
 355 360 365
 Pro Val Ile Glu Ser Leu Gly Leu Pro Gly Phe Gly Tyr His Ser Asp
 370 375 380
 25 Lys Ala Glu Tyr Val Asp Ile Ser Ala Ile Pro Arg Arg Leu Tyr Met
 385 390 395 400
 Ala Ala Arg Leu Ile Met Asp Leu Gly Ala Gly Lys
 405 410

30 (2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

CCACTCTCAC AGTGAGCTCG G

21

40

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 55 base pairs

(B) TYPE: nucleic acid

45 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- 67 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

ACCGCTACCG CCACCACCAG AGCCACCACC GCCAACTGTC TTGTCCACCT TGGTG 55

5 (2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

ACCCCCTCTA GAGTCGAC 18

15

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 54 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

25 TCTGGTGGTG GCGGTAGCGG TGGCGGGGGT TCCCAGAAGC GCGACAACGT GCTG 54

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1929 base pairs

30 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

35

ATGGAGTTGT GGCTGAACTG GATTTTCCTT GTAACACTTT TAAATGGTAT CCAGTGTGAG	60
GTGAAGCTGG TGGAGTCTGG AGGAGGCTTG GTACAGCCTG GGGGTTCTCT GAGACTCTCC	120
TGTGCAACTT CTGGGTTTAC CTTCACTGAT TACTACATGA ACTGGGTCCG CCAGCCTCCA	180
GGAAAGGCAC TTGAGTGGTT GGGTTTATT GGAAACAAAG CTAATGGTTA CACAACAGAG	240
40 TACAGTGCAT CTGTGAAGGG TCGGTTACAC ATCTCCAGAG ATAAATCCCA AAGCATCCTC	300
TATCTTCAAA TGAACACCCT GAGAGCTGAG GACAGTGCCA CTTATTACTG TACAAGAGAT	360
AGGGGGCTAC GGTTCTACTT TGA TACTACTGG GGCCAAGGCA CCACTCTCAC AGTGAGCTCG	420
GCTAGCACCA AGGGACCATC GGTCTTCCCC CTGGCCCCCT GCTCCAGGAG CACCTCCGAG	480
AGCACAGCCG CCCTGGGCTG CCTGGTCAAG GACTACTTCC CCGAACCAGG GACGGGTGTCG	540
45 TGGAAGCTCAG GCGCTCTGAC CAGCGGCGTG CACACCTTCC CGGCTGTCCT ACAGTCCTCA	600
GGACTCTACT CCCTCAGCAG CGTCGTGACG GTGCCCTCCA GCAACTTCGG CACCCAGACC	660
TACACCTGCA ACGTAGATCA CAAGCCCAGC AACACCAAGG TGGACAAGAC AGTTGGCGGT	720

- 68 -

GTGGGCTCTG GTGGTGGCGG TAGCGGTGGC GGGGGTTCCC AGAAGCGCGA CAACGTGCTG 780
 TTCCAGGCAG CTACCGACGA GCAGCCGGCC GTGATCAAGA CGCTGGAGAA GCTGGTCAAC 840
 ATCGAGACCG GCACCGGTGA CGCCGAGGGC ATCGCCGCTG CGGGCAACTT CCTCGAGGCC 900
 GAGCTCAAGA ACCTCGGCTT CACGGTCACG CGAAGCAAGT CGGCCGGCCT GGTGGTGGGC 960
 5 GACAAACATCG TGGGCAAGAT CAAGGGCCGC GGCAGCAAGA ACCTGCTGCT GATGTCGCAC 1020
 ATGGACACCG TCTACCTCAA GGGCATTCTC GCGAAGGCCC CGTTCCGCGT CGAAGGCGAC 1080
 AAGGCCTACG GCCCGGGCAT CGCCGACGAC AAGGGCGGCA ACGCGGTCAT CCTGCACACG 1140
 CTCAGCTGC TGAAGGAATA CGGCGTGC GCCTACGGCA CCATCACCCT GCTGTTCAAC 1200
 ACCGACGAGG AAAAGGGTTC CTTCGGCTCG CGCGACCTGA TCCAGGAAGA AGCCAAGCTG 1260
 10 GCCGACTACG TGCTCTCCTT CGAGCCCACC AGCGCAGGCG ACGAAAACT CTCGCTGGGC 1320
 ACCTCGGGCA TGCCTACGT GCAGGTCCAG ATCACCAGCA AGGCCTCGCA TGCCGGCGCC 1380
 GCGCCCGAGC TGGCGTGAA CGCGCTGGTC GAGGCTTCCG ACCTCGTGCT GCGCACGATG 1440
 AACATCGACG ACAAGGCGAA GAACCTGCGC TTCCAGTGGA CCATCGCCAA GGCCGGCCAG 1500
 GTCTCGAACA TCATCCCCGC CAGCGCCACG CTGAACGCGC ACGTGCCTA CGCGCGCAAC 1560
 15 GAGGACTTCG ACGCCGCCAT GAAGACGCTG GAAGAGCGCG CGCAGCAGAA GAAGCTGCCC 1620
 GAGGCCGACG TGAAGGTGAT CGTCACGCGC GGC CGCCCGG CCTTCAATGC CGGCGAAGGC 1680
 GGCAAGAAGC TGGTCGACAA GGCGGTGGCC TACTACAAGG AAGCCGGCGG CACGCTGGGC 1740
 GTGAAGAGC GCACCGGCGG CGGCACCGAC GCGGCCTACG CCGCGCTCTC AGGCAAGCCA 1800
 GTGATCGAGA GCCTGGGCCT GCCGGGCTTC GGCTACCACA GCGACAAGGC CGAGTACGTG 1860
 20 GACATCAGCG CGATTCCGCG CCGCTGTAC ATGGCTGCGC GCCTGATCAT GGATCTGGGC 1920
 GCCGGCAAG 1929

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 643 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Met Glu Leu Trp Leu Asn Trp Ile Phe Leu Val Thr Leu Leu Asn Gly
 1 5 10 15
 Ile Gln Cys Glu Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Thr Ser Gly Phe Thr Phe
 35 40 45
 Thr Asp Tyr Tyr Met Asn Trp Val Arg Gln Pro Pro Gly Lys Ala Leu
 50 55 60
 40 Glu Trp Leu Gly Phe Ile Gly Asn Lys Ala Asn Gly Tyr Thr Thr Glu
 65 70 75 80
 Tyr Ser Ala Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Lys Ser
 85 90 95
 Gln Ser Ile Leu Tyr Leu Gln Met Asn Thr Leu Arg Ala Glu Asp Ser
 100 105 110
 45 Ala Thr Tyr Tyr Cys Thr Arg Asp Arg Gly Leu Arg Phe Tyr Phe Asp
 115 120 125

- 69 -

Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Ser Thr Lys
 130 135 140
 Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
 145 150 155 160
 5 Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
 165 170 175
 Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
 180 185 190
 Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
 195 200 205
 10 Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn
 210 215 220
 Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Gly Gly
 225 230 235 240
 15 Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Lys Arg
 245 250 255
 Asp Asn Val Leu Phe Gln Ala Ala Thr Asp Glu Gln Pro Ala Val Ile
 260 265 270
 Lys Thr Leu Glu Lys Leu Val Asn Ile Glu Thr Gly Thr Gly Asp Ala
 275 280 285
 20 Glu Gly Ile Ala Ala Ala Gly Asn Phe Leu Glu Ala Glu Leu Lys Asn
 290 295 300
 Leu Gly Phe Thr Val Thr Arg Ser Lys Ser Ala Gly Leu Val Val Gly
 305 310 315 320
 25 Asp Asn Ile Val Gly Lys Ile Lys Gly Arg Gly Gly Lys Asn Leu Leu
 325 330 335
 Leu Met Ser His Met Asp Thr Val Tyr Leu Lys Gly Ile Leu Ala Lys
 340 345 350
 Ala Pro Phe Arg Val Glu Gly Asp Lys Ala Tyr Gly Pro Gly Ile Ala
 355 360 365
 30 Asp Asp Lys Gly Gly Asn Ala Val Ile Leu His Thr Leu Lys Leu Leu
 370 375 380
 Lys Glu Tyr Gly Val Arg Asp Tyr Gly Thr Ile Thr Val Leu Phe Asn
 385 390 395 400
 35 Thr Asp Glu Glu Lys Gly Ser Phe Gly Ser Arg Asp Leu Ile Gln Glu
 405 410 415
 Glu Ala Lys Leu Ala Asp Tyr Val Leu Ser Phe Glu Pro Thr Ser Ala
 420 425 430
 Gly Asp Glu Lys Leu Ser Leu Gly Thr Ser Gly Ile Ala Tyr Val Gln
 435 440 445
 40 Val Gln Ile Thr Gly Lys Ala Ser His Ala Gly Ala Ala Pro Glu Leu
 450 455 460
 Gly Val Asn Ala Leu Val Glu Ala Ser Asp Leu Val Leu Arg Thr Met
 465 470 475 480
 45 Asn Ile Asp Asp Lys Ala Lys Asn Leu Arg Phe Gln Trp Thr Ile Ala
 485 490 495
 Lys Ala Gly Gln Val Ser Asn Ile Ile Pro Ala Ser Ala Thr Leu Asn

- 70 -

500 505 510
 Ala Asp Val Arg Tyr Ala Arg Asn Glu Asp Phe Asp Ala Ala Met Lys
 515 520 525
 Thr Leu Glu Glu Arg Ala Gln Gln Lys Lys Leu Pro Glu Ala Asp Val
 530 535 540
 Lys Val Ile Val Thr Arg Gly Arg Pro Ala Phe Asn Ala Gly Glu Gly
 545 550 555 560
 Gly Lys Lys Leu Val Asp Lys Ala Val Ala Tyr Tyr Lys Glu Ala Gly
 565 570 575
 10 Gly Thr Leu Gly Val Glu Glu Arg Thr Gly Gly Gly Thr Asp Ala Ala
 580 585 590
 Tyr Ala Ala Leu Ser Gly Lys Pro Val Ile Glu Ser Leu Gly Leu Pro
 595 600 605
 Gly Phe Gly Tyr His Ser Asp Lys Ala Glu Tyr Val Asp Ile Ser Ala
 15 610 615 620
 Ile Pro Arg Arg Leu Tyr Met Ala Ala Arg Leu Ile Met Asp Leu Gly
 625 630 635 640
 Ala Gly Lys

20

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 705 base pairs

(B) TYPE: nucleic acid

25

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

30 ATGGATTTC AAGTGCAGAT TTTCAGCTTC CTGCTAATCA GTGCTTCAGT CATAATGTCC 60
 AGAGGACAAA CTGTTCTCTC CCAGTCTCCA GCAATCCTGT CTGCATCTCC AGGGGAGAAG 120
 GTCACAATGA CTGTCAGGGC CAGCTCAAGT GTAACTTACA TTCACTGGTA CCAGCAGAAG 180
 CCAGGTTCTT CCCCCAAATC CTGGATTTAT GCCACATCCA ACCTGGCTTC TGGAGTCCCT 240
 GCTCGCTTCA GTGGCAGTGG GTCTGGGACC TCTTACTCTC TCACAATCAG CAGAGTGGAG 300
 35 GCTGAAGATG CTGCCACTTA TTAAGTCCAA CATTGGAGTA GTAAACCACC GACGTTCCGT 360
 GGAGGCACCA AGCTCGAGAT CAAACGGACT GTGGCTGCAC CATCTGTCTT CATCTTCCCG 420
 CCATCTGATG AGCAGTTGAA ATCTGGAAGT GCCTCTGTTG TGTGCTGCT GAATAACTTC 480
 TATCCCAGAG AGGCCAAAGT ACAGTGAAG GTGGATAACG CCCTCCAATC GGGTAACTCC 540
 CAGGAGAGTG TCACAGAGCA GGACAGCAAG GACAGCACCT ACAGCCTCAG CAGCACCTG 600
 40 ACGCTGAGCA AAGCAGACTA CGAGAAACAC AAAGTCTACG CCTGCGAAGT CACCCATCAG 660
 GCCTGAGTT CGCCGTCAC AAAGAGCTTC AACAGGGGAG AGTGT 705

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

45

(A) LENGTH: 235 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

- 71 -

- (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: protein
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

```

5   Met Asp Phe Gln Val Gln Ile Phe Ser Phe Leu Leu Ile Ser Ala Ser
    1             5             10             15
    Val Ile Met Ser Arg Gly Gln Thr Val Leu Ser Gln Ser Pro Ala Ile
          20             25             30
    Leu Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
10   35             40             45
    Ser Ser Val Thr Tyr Ile His Trp Tyr Gln Gln Lys Pro Gly Ser Ser
    50             55             60
    Pro Lys Ser Trp Ile Tyr Ala Thr Ser Asn Leu Ala Ser Gly Val Pro
    65             70             75             80
15   Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
          85             90             95
    Ser Arg Val Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Trp
          100            105            110
    Ser Ser Lys Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
20   115            120            125
    Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
          130            135            140
    Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
    145            150            155            160
25   Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
          165            170            175
    Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
          180            185            190
    Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
30   195            200            205
    Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
          210            215            220
    Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
    225            230            235
35

```

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39 base pairs

(B) TYPE: nucleic acid

40 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

45 AAGCTTGAAT TCGCCGCCAC TATGGATTTT CAAGTGCAG

39

(2) INFORMATION FOR SEQ ID NO: 20:

- 72 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 44 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
5 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: other nucleic acid
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:
- TTAATTGGAT CCGAGCTCCT ATTAACACTC TCCCCTGTTG AAGC 44
- 10
- (2) INFORMATION FOR SEQ ID NO: 21:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 50 base pairs
 (B) TYPE: nucleic acid
15 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: other nucleic acid
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:
- 20 AAGCTTCCGG ATCCCTGCAG CCATGGAGTT GTGGCTGAAC TGGATTTTCC 50
- (2) INFORMATION FOR SEQ ID NO: 22:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 base pairs
25 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: other nucleic acid
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:
- 30
- AAGCTTAGTC TAGATTATCA CTTGCCGGCG CCCAGATC 38
- (2) INFORMATION FOR SEQ ID NO: 23:
 (i) SEQUENCE CHARACTERISTICS:
35 (A) LENGTH: 46 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: other nucleic acid
40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:
- CGGGGGATCC AGATCTGAGC TCCTGTAGAC GTCGACATTA ATTCCG 46
- (2) INFORMATION FOR SEQ ID NO: 24:
45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid

- 73 -

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

5

GGAAAATCCA GTTCAGCCAC AACTCCATGG

30

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 1926 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

ATGAAGTTGT GGCTGAAGTG GATTTTCCTT GTAACACTTT TAAATGGAAT TCAGTGTGAG	60
GTGCAGCTGC AGCAGTCTGG GGCAGAGCTT GTGAGGTCAG GGGCCTCAGT CAAGTTGTCC	120
TGCACAGCTT CTGGCTTCAA CATTAAAGAC AACTATATGC ACTGGGTGAA GCAGAGGCCT	180
20 GAACAGGGCC TGGAGTGGAT TGCATGGATT GATCCTGAGA ATGGTGATAC TGAATATGCC	240
CCGAAGTTCC GGGGCAAGGC CACTTTGACT GCAGACTCAT CCTCCAACAC AGCCTACCTG	300
CACCTCAGCA GCCTGACATC TGAGGACACT GCCGTCTATT ACTGTCATGT CCTGATCTAT	360
GCTGGTTATT TGGCTATGGA CTACTGGGGT CAAGGAACCT CAGTCGCCGT GAGCTCGGCT	420
AGCACCAAGG GACCATCGGT CTTCCTCCCTG GCCCCCTGCT CCAGGAGCAC CTCCGAGAGC	480
25 ACAGCCGCCC TGGGCTGCCT GGTCAAGGAC TACTTCCCCG AACCGGTGAC GGTGTCTGTTG	540
AACTCAGGCG CTCTGACCAG CGGCGTGAC ACCTTCCCGG CTGTCTTACA GTCTCAGGA	600
CTCTACTCCC TCAGCAGCGT CGTGACGGTG CCCTCCAGCA ACTTCGGCAC CCAGACCTAC	660
ACCTGCAACG TAGATCACAA GCCCAGCAAC ACCAAGGTGG ACAAGACAGT TGGCGGTGGT	720
GGCTCTGGTG GTGGCGGTAG CGGTGGCGGG GGTTCCTCAGA AGCGCGACAA CGTGCTGTTC	780
30 CAGGCAGCTA CCGACGAGCA GCCGGCCGTG ATCAAGACGC TGGAGAAGCT GGTCAACATC	840
GAGACCGGCA CCGGTGACGC CGAGGGCATC GCCGCTGCGG GCAACTTCCT CGAGGCCGAG	900
CTCAAGAACC TCGGCTTCAC GGTACGCGA AGCAAGTCGG CCGGCCTGGT GGTGGGCGAC	960
AACATCGTGG GCAAGATCAA GGGCCGCGGC GGCAAGAACC TGCTGCTGAT GTCGCACATG	1020
GACACCGTCT ACCTCAAGGG CATTCTCGCG AAGGCCCGGT TCCGCGTCGA AGGCGACAAG	1080
35 GCCTACGGCC CGGGCATCGC CGACGACAAG GCGGCAACG CGGTCATCCT GCACACGCTC	1140
AAGCTGCTGA AGGAATACGG CGTGCGCGAC TACGGACCA TCACCGTGCT GTTCAACACC	1200
GACGAGGAAA AGGGTTCCTT CGGCTCGCGC GACCTGATCC AGGAAGAAGC CAAGCTGGCC	1260
GACTACGTGC TCTCCTTCGA GCCCACCAGC GCAGGCGACG AAAAACTCTC GCTGGGCACC	1320
TCGGGCATCG CCTACGTGCA GGTCCAGATC ACCGGCAAGG CCTCGCATGC CGGCGCCGCG	1380
40 CCCGAGCTGG GCGTGAACGC GCTGGTCGAG GCTTCCGACC TCGTGCTGCG CACGATGAAC	1440
ATCGACGACA AGGCGAAGAA CCTGCGCTTC CAGTGGACCA TCGCCAAGGC CGGCCAGGTC	1500
TGGAACATCA TCCCGCCAG CGCCACGCTG AACCGCGACG TCGCTACGC GCGCAACGAG	1560
GACTTCGACG CCGCATGAA GACGCTGGAA GAGCGCGCGC AGCAGAAGAA GCTGCCCGAG	1620
GCCGACGTGA AGGTGATCGT CACGCGCGGC CGCCCGGCCT TCAATGCCGG CGAAGGCGGC	1680

- 74 -

AAGAAGCTGG TCGACAAGGC GGTGGCCTAC TACAAGGAAG CCGGCGGCAC GCTGGGCGTG 1740
 GAAGAGCGCA CCGGCGGCGG CACCGACGCG GCCTACGCCG CGCTCTCAGG CAAGCCAGTG 1800
 ATCGAGAGCC TGGGCTTGGC GGGCTTCGGC TACCACAGCG ACAAGGCCGA GTACGTGGAC 1860
 ATCAGCGCGA TTCCGCGCGC CCTGTACATG GCTGCGCGCC TGATCATGGA TCTGGGCGCC 1920
 5 GGCAAG 1926

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 642 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

15

Met Lys Leu Trp Leu Asn Trp Ile Phe Leu Val Thr Leu Leu Asn Gly
 1 5 10 15
 Ile Gln Cys Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg
 20 25 30
 Ser Gly Ala Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile
 35 40 45
 Lys Asp Asn Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu
 50 55 60
 Glu Trp Ile Ala Trp Ile Asp Pro Glu Asn Gly Asp Thr Glu Tyr Ala
 25 65 70 75 80
 Pro Lys Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Ser Ser Ser Asn
 85 90 95
 Thr Ala Tyr Leu His Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys His Val Leu Ile Tyr Ala Gly Tyr Leu Ala Met Asp Tyr
 115 120 125
 Trp Gly Gln Gly Thr Ser Val Ala Val Ser Ser Ala Ser Thr Lys Gly
 130 135 140
 Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser
 145 150 155 160
 Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val
 165 170 175
 Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe
 180 185 190
 Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val
 195 200 205
 Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn Val
 210 215 220
 Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Gly Gly Gly
 45 225 230 235 240
 Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Lys Arg Asp
 245 250 255

- 75 -

Asn Val Leu Phe Gln Ala Ala Thr Asp Glu Gln Pro Ala Val Ile Lys
 260 265 270
 Thr Leu Glu Lys Leu Val Asn Ile Glu Thr Gly Thr Gly Asp Ala Glu
 275 280 285
 5 Gly Ile Ala Ala Ala Gly Asn Phe Leu Glu Ala Glu Leu Lys Asn Leu
 290 295 300
 Gly Phe Thr Val Thr Arg Ser Lys Ser Ala Gly Leu Val Val Gly Asp
 305 310 315 320
 Asn Ile Val Gly Lys Ile Lys Gly Arg Gly Gly Lys Asn Leu Leu Leu
 10 325 330 335
 Met Ser His Met Asp Thr Val Tyr Leu Lys Gly Ile Leu Ala Lys Ala
 340 345 350
 Pro Phe Arg Val Glu Gly Asp Lys Ala Tyr Gly Pro Gly Ile Ala Asp
 355 360 365
 15 Asp Lys Gly Gly Asn Ala Val Ile Leu His Thr Leu Lys Leu Leu Lys
 370 375 380
 Glu Tyr Gly Val Arg Asp Tyr Gly Thr Ile Thr Val Leu Phe Asn Thr
 385 390 395 400
 Asp Glu Glu Lys Gly Ser Phe Gly Ser Arg Asp Leu Ile Gln Glu Glu
 20 405 410 415
 Ala Lys Leu Ala Asp Tyr Val Leu Ser Phe Glu Pro Thr Ser Ala Gly
 420 425 430
 Asp Glu Lys Leu Ser Leu Gly Thr Ser Gly Ile Ala Tyr Val Gln Val
 435 440 445
 25 Gln Ile Thr Gly Lys Ala Ser His Ala Gly Ala Ala Pro Glu Leu Gly
 450 455 460
 Val Asn Ala Leu Val Glu Ala Ser Asp Leu Val Leu Arg Thr Met Asn
 465 470 475 480
 Ile Asp Asp Lys Ala Lys Asn Leu Arg Phe Gln Trp Thr Ile Ala Lys
 30 485 490 495
 Ala Gly Gln Val Ser Asn Ile Ile Pro Ala Ser Ala Thr Leu Asn Ala
 500 505 510
 Asp Val Arg Tyr Ala Arg Asn Glu Asp Phe Asp Ala Ala Met Lys Thr
 515 520 525
 35 Leu Glu Glu Arg Ala Gln Gln Lys Lys Leu Pro Glu Ala Asp Val Lys
 530 535 540
 Val Ile Val Thr Arg Gly Arg Pro Ala Phe Asn Ala Gly Glu Gly Gly
 545 550 555 560
 Lys Lys Leu Val Asp Lys Ala Val Ala Tyr Tyr Lys Glu Ala Gly Gly
 40 565 570 575
 Thr Leu Gly Val Glu Glu Arg Thr Gly Gly Gly Thr Asp Ala Ala Tyr
 580 585 590
 Ala Ala Leu Ser Gly Lys Pro Val Ile Glu Ser Leu Gly Leu Pro Gly
 595 600 605
 45 Phe Gly Tyr His Ser Asp Lys Ala Glu Tyr Val Asp Ile Ser Ala Ile
 610 615 620
 Pro Arg Arg Leu Tyr Met Ala Ala Arg Leu Ile Met Asp Leu Gly Ala

- 76 -

625

630

635

640

Gly Lys

- 5 (2) INFORMATION FOR SEQ ID NO: 27:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - 10 (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: other nucleic acid
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

AAGCTTGGAA TTCAGTGTCA GGTCCAACCTG CAGCAGCCT

39

15

- (2) INFORMATION FOR SEQ ID NO: 28:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 54 base pairs
 - (B) TYPE: nucleic acid
 - 20 (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: other nucleic acid
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

25 GCTACCGCCA CCTCCGGAGC CACCACCGCC CCGTTTGATC TCGAGCTTGG TGCC

54

- (2) INFORMATION FOR SEQ ID NO: 29:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 58 base pairs
 - 30 (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: other nucleic acid
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

35

TCCGGAGGTG GCGGTAGCGG TGGCGGGGGT TCCAGAAGC GCGACAACGT GCTGTTCC

58

- (2) INFORMATION FOR SEQ ID NO: 30:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - 40 (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: other nucleic acid
 - 45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

CCTCGAGGAA TTCTTCACT TGCC

24

- 77 -

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 2019 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

10
 ATGAAGTTGT GGCTGAAC TG GATTTTCCTT GTAACACTTT TAAATGGAAT TCAGTGTGTCAG 60
 GTCCAACTGC AGCAGCCTGG GGCTGAAC TG GTGAAGCCTG GGGCTTCAGT GCAGCTGTCC 120
 TGCAAGGCTT CTGGCTACAC CTTCAACGGC TACTGGATAC ACTGGGTGAA GCAGAGGCCT 180
 GGACAAGGCC TTGAGTGGAT TGGAGAGGTT AATCCTAGTA CCGGTCGTTC TGACTACAAT 240
 15 GAGAAGTTCA AGAACAAGGC CAACTGACT GTAGACAAAT CCTCCACCAC AGCCTACATG 300
 CAACTCAGCA GCCTGACATC TGAGGACTCT GCGGTCTATT ACTGTGCAAG AGAGAGGGCC 360
 TATGGTTACG ACGATGCTAT GGACTACTGG GGCCAAGGGA CCACGGTCAC CGTCTCCTCA 420
 GGTGGCGGTG GCTCGGGCGG TGGTGGGTCG GGTGGCGGCG GATCTGACAT TGAGCTCTCA 480
 CAGTCTCCAT CCTCCCTGGC TGTGTCAGCA GGAGAGAAGG TCACCATGAG CTGCAAATCC 540
 20 AGTCAGAGTC TCCTCAACAG TAGAACCCGA AAGAACTACT TGGCTTGTA CCAGCAGAGA 600
 CCAGGGCAGT CTCCTAACT GCTGATCTAT TGGGCATCCA CTAGGACATC TGGGGTCCCT 660
 GATCGCTTCA CAGGCAGTGG ATCTGGGACA GATTTCACTC TCACCATCAG CAGTGTGCAG 720
 GCTGAAGACC TGGCAATTTA TTAAGTCAAG CAATCTTATA CTCTTCGGAC GTTCGGTGGA 780
 GGCACCAAGC TCGAGATCAA ACGGGGCGGT GGTGGCTCCG GAGGTGGCGG TAGCGGTGGC 840
 25 GGGGGTTCCC AGAAGCGCGA CAACGTGCTG TTCCAGGCAG CTACCGACGA GCAGCCGGCC 900
 GTGATCAAGA CGTGGAGAA GCTGGTCAAC ATCGAGACCG GCACCGGTGA CGCCGAGGGC 960
 ATCGCCGCTG CGGGCAACTT CCTCGAGGCC GAGCTCAAGA ACCTCGGCTT CACGGTCACG 1020
 CGAAGCAAGT CGGCCGGCCT GGTGGTGGGC GACAACATCG TGGGCAAGAT CAAGGGCCGC 1080
 GCGGCAAGA ACCTGCTGCT GATGTCGCAC ATGGACACCG TCTACCTCAA GGGCATTCTC 1140
 30 GCGAAGGCCC CGTCCGCGT CGAAGGCGAC AAGGCCTACG GCGCGGCAT CGCCGACGAC 1200
 AAGGGCGGCA ACGCGGTCAT CCTGCACACG CTCAAGCTGC TGAAGGAATA CGGCGTGGC 1260
 GACTACGGCA CCATACCGT GCTGTTCAAC ACCGACGAGG AAAAGGGTTC CTTGCGCTCG 1320
 CGCGACCTGA TCCAGGAAGA AGCCAAGCTG GCCGACTACG TGCTCTCCTT CGAGCCCACC 1380
 AGCGCAGGCG ACGAAAACT CTCGCTGGGC ACCTCGGGCA TCGCCTACGT CGAGGTCCAG 1440
 35 ATCACCGGCA AGGCCTCGCA TGCCGGCGCC GCGCCGAGC TGGCGTGAA CGCGCTGGTC 1500
 GAGGCTTCCG ACCTCGTGCT GCGCACGATG AACATCGAG ACAAGGCGAA GAACCTGCGC 1560
 TTCCAGTGGA CCATCGCCAA GGCCGGCCAG GTCTCGAACA TCATCCCCGC CAGCGCCACG 1620
 CTGAACGCCG ACGTGCCTA CGCGCGCAAC GAGGACTTCG ACGCCGCCAT GAAGACGCTG 1680
 GAAGAGCGCG CGCAGCAGAA GAAGCTGCCC GAGGCCGAG TGAAGGTGAT CGTCACGCGC 1740
 40 GGCCGCCCGG CTTCAATGC CGGCGAAGGC GGCAAGAAGC TGSTCGCAA GGCGGTGGCC 1800
 TACTACAAGG AAGCCGGCGG CACGCTGGGC GTGGAAGAGC GCACCGGCGG CGGCACCGAC 1860
 GCGGCCTACG CCGCGCTCTC AGGCAAGCCA GTGATCGAGA GCCTGGGCTT GCCGGGCTTC 1920
 GGCTACCACA GCGACAAGGC CGAGTACGTG GACATCAGCG CGATTCCGCG CCGCTGTAC 1980
 ATGGCTGCGC GCCTGATCAT GGATCTGGGC GCCGGCAAG 2019

45

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:

- 78 -

(A) LENGTH: 673 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

```

Met Lys Leu Trp Leu Asn Trp Ile Phe Leu Val Thr Leu Leu Asn Gly
 1             5             10             15
10 Ile Gln Cys Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys
    20             25             30
Pro Gly Ala Ser Val Gln Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe
    35             40             45
15 Thr Gly Tyr Trp Ile His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu
    50             55             60
Glu Trp Ile Gly Glu Val Asn Pro Ser Thr Gly Arg Ser Asp Tyr Asn
65             70             75             80
Glu Lys Phe Lys Asn Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Thr
    85             90             95
20 Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val
    100            105            110
Tyr Tyr Cys Ala Arg Glu Arg Ala Tyr Gly Tyr Asp Asp Ala Met Asp
    115            120            125
25 Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly Gly Gly
    130            135            140
Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile Glu Leu Ser
145            150            155            160
Gln Ser Pro Ser Ser Leu Ala Val Ser Ala Gly Glu Lys Val Thr Met
    165            170            175
30 Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser Arg Thr Arg Lys Asn
    180            185            190
Tyr Leu Ala Trp Tyr Gln Gln Arg Pro Gly Gln Ser Pro Lys Leu Leu
    195            200            205
35 Ile Tyr Trp Ala Ser Thr Arg Thr Ser Gly Val Pro Asp Arg Phe Thr
    210            215            220
Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Val Gln
225            230            235            240
Ala Glu Asp Leu Ala Ile Tyr Tyr Cys Lys Gln Ser Tyr Thr Leu Arg
    245            250            255
40 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Gly Gly Gly Gly
    260            265            270
Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Lys Arg Asp Asn
    275            280            285
45 Val Leu Phe Gln Ala Ala Thr Asp Glu Gln Pro Ala Val Ile Lys Thr
    290            295            300
Leu Glu Lys Leu Val Asn Ile Glu Thr Gly Thr Gly Asp Ala Glu Gly
305            310            315            320

```

- 79 -

Ile Ala Ala Ala Gly Asn Phe Leu Glu Ala Glu Leu Lys Asn Leu Gly
 325 330 335
 Phe Thr Val Thr Arg Ser Lys Ser Ala Gly Leu Val Val Gly Asp Asn
 340 345 350
 5 Ile Val Gly Lys Ile Lys Gly Arg Gly Gly Lys Asn Leu Leu Leu Met
 355 360 365
 Ser His Met Asp Thr Val Tyr Leu Lys Gly Ile Leu Ala Lys Ala Pro
 370 375 380
 10 Phe Arg Val Glu Gly Asp Lys Ala Tyr Gly Pro Gly Ile Ala Asp Asp
 385 390 395 400
 Lys Gly Gly Asn Ala Val Ile Leu His Thr Leu Lys Leu Leu Lys Glu
 405 410 415
 Tyr Gly Val Arg Asp Tyr Gly Thr Ile Thr Val Leu Phe Asn Thr Asp
 420 425 430
 15 Glu Glu Lys Gly Ser Phe Gly Ser Arg Asp Leu Ile Gln Glu Glu Ala
 435 440 445
 Lys Leu Ala Asp Tyr Val Leu Ser Phe Glu Pro Thr Ser Ala Gly Asp
 450 455 460
 20 Glu Lys Leu Ser Leu Gly Thr Ser Gly Ile Ala Tyr Val Gln Val Gln
 465 470 475 480
 Ile Thr Gly Lys Ala Ser His Ala Gly Ala Ala Pro Glu Leu Gly Val
 485 490 495
 Asn Ala Leu Val Glu Ala Ser Asp Leu Val Leu Arg Thr Met Asn Ile
 500 505 510
 25 Asp Asp Lys Ala Lys Asn Leu Arg Phe Gln Trp Thr Ile Ala Lys Ala
 515 520 525
 Gly Gln Val Ser Asn Ile Ile Pro Ala Ser Ala Thr Leu Asn Ala Asp
 530 535 540
 30 Val Arg Tyr Ala Arg Asn Glu Asp Phe Asp Ala Ala Met Lys Thr Leu
 545 550 555 560
 Glu Glu Arg Ala Gln Gln Lys Lys Leu Pro Glu Ala Asp Val Lys Val
 565 570 575
 Ile Val Thr Arg Gly Arg Pro Ala Phe Asn Ala Gly Glu Gly Gly Lys
 580 585 590
 35 Lys Leu Val Asp Lys Ala Val Ala Tyr Tyr Lys Glu Ala Gly Gly Thr
 595 600 605
 Leu Gly Val Glu Glu Arg Thr Gly Gly Gly Thr Asp Ala Ala Tyr Ala
 610 615 620
 40 Ala Leu Ser Gly Lys Pro Val Ile Glu Ser Leu Gly Leu Pro Gly Phe
 625 630 635 640
 Gly Tyr His Ser Asp Lys Ala Glu Tyr Val Asp Ile Ser Ala Ile Pro
 645 650 655
 Arg Arg Leu Tyr Met Ala Ala Arg Leu Ile Met Asp Leu Gly Ala Gly
 660 665 670
 45 Lys

- 80 -

- (2) INFORMATION FOR SEQ ID NO: 33:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 37 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:
- 10 GGGCGCCGGC AAGTGATAAA ATTCCTCGAG GAGCTCC 37
- (2) INFORMATION FOR SEQ ID NO: 34:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:
- 15 CGCCACCTCT GACTTGAGC 19
- (2) INFORMATION FOR SEQ ID NO: 35:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 37 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:
- 25 30 GGAGCTCCTC GAGGAATTTT ATCACTTGCC GCGCCCC 37
- (2) INFORMATION FOR SEQ ID NO: 36:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:
- 35 40 GCTGAACGCC GACGTGCGC 19
- 45 (2) INFORMATION FOR SEQ ID NO: 37:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2025 base pairs

- 81 -

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

```

ATGAAGTTGT GGCTGAACTG GATTTTCCTT GTAACACTTT TAAATGGAAT TCAGTGTCTCAG      60
GTCCAACCTGC AGCAGCCTGG GGCTGAACTG GTGAAGCCTG GGGCTTCAGT GCAGCTGTCC      120
TGCAAGGCTT CTGGCTACAC CTTCAACCGGC TACTGGATAC ACTGGGTGAA GCAGAGGCCT      180
10 GGACAAGGCC TTGAGTGGAT TGGAGAGGTT AATCCTAGTA CCGGTCGTTC TGACTACAAT      240
GAGAAGTTCA AGAACAAGGC CACACTGACT GTAGACAAAT CCTCCACCAC AGCCTACATG      300
CAACTCAGCA GCCTGACATC TGAGGACTCT GCGGTCTATT ACTGTGCAAG AGAGAGGGCC      360
TATGGTTACG ACGATGCTAT GGACTACTGG GGCCAAGGGA CCACGGTCAC CGTCTCCTCA      420
GGTGGCGGTG GCTCGGGCGG TGGTGGGTCG GGTGGCGGCG GATCTGACAT TGAGCTCTCA      480
15 CAGTCTCCAT CCTCCCTGGC TGTGTCTAGCA GGAGAGAAGG TCACCATGAG CTGCAAATCC      540
AGTCAGAGTC TCCTCAACAG TAGAACCCGA AAGAACTACT TGGCTTGGTA CCAGCAGAGA      600
CCAGGGCAGT CTCCTAAACT GCTGATCTAT TGGGCATCCA CTAGGACATC TGGGGTCCCT      660
GATCGCTTCA CAGGCAGTGG ATCTGGGACA GATTTCACTC TCACCATCAG CAGTGTGCAG      720
GCTGAAGACC TGGCAATTTA TTACTGCAAG CAATCTTATA CTCTTCGGAC GTTCGGTGGA      780
20 GGCACCAAGC TCGAGATCAA ACGGGGCGGT GGTGGCTCCG GAGGTGGCGG TAGCCGTGGC      840
GGGGGTTCCT AGAAGCGCGA CAACGTGCTG TTCCAGGCAG CTACCGACGA GCAGCCGGCC      900
GTGATCAAGA CGCTGGAGAA GCTGGTCAAC ATCGAGACCG GCACCGGTGA CGCCGAGGGC      960
ATCGCCGCTG CGGGCAACTT CCTCGAGGCC GAGCTCAAGA ACCTCGGCTT CACGGTCACG      1020
CGAAGCAAGT CGGCCGGCCT GGTGGTGGGC GACAACATCG TGGGCAAGAT CAAGGGCCGC      1080
25 GCGCGCAAGA ACCTGCTGCT GATGTCGCAC ATGGACACCG TCTACCTCAA GGGCATTCTC      1140
GCGAAGGCCC CGTTCCGCGT CGAAGGCGAC AAGGCCTACG GCGCGGCAT CGCCGACGAC      1200
AAGGGCGGCA ACGCGGTCAT CCTGCACACG CTCAAGCTGC TGAAGGAATA CGGCGTGCGC      1260
GACTACGGCA CCATCACCCT GCTGTTCAAC ACCGACGAGG AAAAGGGTTC CTTCGGCTCG      1320
CGCGACCTGA TCCAGGAAGA AGCCAAGCTG GCCGACTACG TGCTCTCCTT CGAGCCCACC      1380
30 AGCGCAGGCG ACGAAAACT CTCGTGGGC ACCTCGGGCA TCGCCTACGT GCAGGTCCAG      1440
ATCACCGGCA AGGCCTCGCA TGCCGGCGCC GCGCCGAGC TGGGCGTGAA CGCGCTGGTC      1500
GAGGCTTCCG ACCTCGTGCT GCGCACGATG AACATCGACG ACAAGGCGAA GAACCTGCGC      1560
TTCCAGTGGA CCATCGCCAA GGCCGGCCAG GTCTCGAACA TCATCCCCGC CAGCGCCACG      1620
CTGAACGCCG ACGTGCCTA CGCGCGCAAC GAGGACTTCG ACGCCGCCAT GAAGACGCTG      1680
35 GAAGAGCGCG CGCAGCAGAA GAAGCTGCCC GAGGCCGACG TGAAGGTGAT CGTCACGCGC      1740
GGCCGCCCGG CTTCAATGC CGGCGAAGGC GGCAAGAAGC TGGTGACAAA GCGGCTGGCC      1800
TACTACAAGG AAGCCGGCGG CACGCTGGGC GTGGAAGAGC GCACCGGCGG CGGCACCGAC      1860
GCGGCCTACG CCGCGCTCTC AGGCAAGCCA GTGATCGAGA GCCTGGGCTT GCCGGGCTTC      1920
GGCTACCACA GCGACAAGGC CGAGTACGTG GACATCAGCG CGATTCCGCG CCGCCTGTAC      1980
40 ATGGCTGCGC GCCTGATCAT GGATCTGGGC GCCGGCAAGT GATAA      2025

```

(2) INFORMATION FOR SEQ ID NO: 38:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 288 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

45

- 82 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

5 Met Lys Tyr Leu Leu Pro Thr Ala Ala Ala Gly Leu Leu Leu Ala
 1 5 10 15
 Ala Gln Pro Ala Met Ala Gln Val Gln Leu Gln Gln Pro Gly Ala Glu
 20 25 30
 Leu Val Lys Pro Gly Ala Ser Val Gln Leu Ser Cys Lys Ala Ser Gly
 35 40 45
 10 Tyr Thr Phe Thr Gly Tyr Trp Ile His Trp Val Lys Gln Arg Pro Gly
 50 55 60
 Gln Gly Leu Glu Trp Ile Gly Glu Val Asn Pro Ser Thr Gly Arg Ser
 65 70 75 80
 Asp Tyr Asn Glu Lys Phe Lys Asn Lys Ala Thr Leu Thr Val Asp Lys
 15 85 90 95
 Ser Ser Thr Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp
 100 105 110
 Ser Ala Val Tyr Tyr Cys Ala Arg Glu Arg Ala Tyr Gly Tyr Asp Asp
 115 120 125
 20 Ala Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly
 130 135 140
 Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile
 145 150 155 160
 Glu Leu Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Ala Gly Glu Lys
 165 170 175
 25 Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser Arg Thr
 180 185 190
 Arg Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Arg Pro Gly Gln Ser Pro
 195 200 205
 30 Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Thr Ser Gly Val Pro Asp
 210 215 220
 Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 225 230 235 240
 Ser Val Gln Ala Glu Asp Leu Ala Ile Tyr Tyr Cys Lys Gln Ser Tyr
 245 250 255
 35 Thr Leu Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Glu
 260 265 270
 Gln Lys Leu Ile Ser Glu Glu Asp Leu Asn His His His His His
 275 280 285

40

(2) INFORMATION FOR SEQ ID NO: 39:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs

(B) TYPE: nucleic acid

45

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- 83 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

GCCCAACCAG CCATGGCCGA GGTGCAGCTG CAGCAG

36

5 (2) INFORMATION FOR SEQ ID NO: 40:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 54 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

CGACCCACCA CCGCCCGAGC CACCGCCACC CGAGCTCAGG GCGACTGAGG TTCC

54

15

(2) INFORMATION FOR SEQ ID NO: 41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 54 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

25 TCGGGCGGTG GTGGGTCGGG TGGCGGCGGA TCTCAGATTG TGCTCACCCA GTCT

54

(2) INFORMATION FOR SEQ ID NO: 42:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

30 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

35

CCGTTTGATC TCGAGCTTGG TCCC

24

(2) INFORMATION FOR SEQ ID NO: 43:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 843 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

ATGAAATACC TATTGCCTAC GGCAGCCGCT GGATTGTTAT TACTCGCTGC CCAACCAGCC

60

- 84 -

ATGGCCGAGG TGCAGCTGCA GCAGTCTGGG GCAGAGCTTG TGAGGTCAGG GGCCTCAGTC 120
 AAGTTGTCCT GCACAGCTTC TGGCTTCAAC ATTAAAGACA ACTATATGCA CTGGGTGAAG 180
 CAGAGGCCCTG AACAGGGCCT GGAGTGGATT GCATGGATTG ATCCTGAGAA TGGTGATACT 240
 GAATATGCCC CGAAGTCCG GGGCAAGGCC ACTTTGACTG CAGACTCATC CTCCAACACA 300
 5 GCCTACCTGC ACCTCAGCAG CCTGACATCT GAGGACACTG CCGTCTATTA CTGTCATGTC 360
 CTGATCTATG CTGTTATTT GGCTATGGAC TACTGGGGTC AAGGAACCTC AGTCGCCGTG 420
 AGCTCGGGTG GCGGTGGCTC GGGCGGTGGT GGGTCGGGTG GCGGCGGATC TCAGATTGTG 480
 CTCACCCACT CTCCAGCAAT CATGTCTGCA TCTCCAGGGG AGAAGGTCAC CATAACCTGC 540
 AGTGCCAGCT CAAGTGTAAC TTACATGCAC TGGTTCAGC AGAAGCCAGG CACTTCTCCC 600
 10 AACTCTGGA TTTATAGCAC ATCCAACCTG GCTTCTGGAG TCCCTGCTCG CTTAGTGGC 660
 AGTGGATCTG GGACCTCTTA CTCTCTCACA ATCAGCCGAA TGGAGGCTGA AGATGCTGCC 720
 ACTTATTACT GCCAGCAAAG GAGTACTTAC CCGCTCACGT TCGGTGCTGG GACCAAGCTC 780
 GAGATCAAAC GGAACAAAA ACTCATCTCA GAAGAAGATC TGAATCACCA CCATCACCAC 840
 CAT 843

15

(2) INFORMATION FOR SEQ ID NO: 44:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 281 amino acids

(B) TYPE: amino acid

20

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

25 Met Lys Tyr Leu Leu Pro Thr Ala Ala Ala Gly Leu Leu Leu Leu Ala
 1 5 10 15
 Ala Gln Pro Ala Met Ala Glu Val Gln Leu Gln Gln Ser Gly Ala Glu
 20 25 30
 30 Leu Val Arg Ser Gly Ala Ser Val Lys Leu Ser Cys Thr Ala Ser Gly
 35 40 45
 Phe Asn Ile Lys Asp Asn Tyr Met His Trp Val Lys Gln Arg Pro Glu
 50 55 60
 Gln Gly Leu Glu Trp Ile Ala Trp Ile Asp Pro Glu Asn Gly Asp Thr
 65 70 75 80
 35 Glu Tyr Ala Pro Lys Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Ser
 85 90 95
 Ser Ser Asn Thr Ala Tyr Leu His Leu Ser Ser Leu Thr Ser Glu Asp
 100 105 110
 40 Thr Ala Val Tyr Tyr Cys His Val Leu Ile Tyr Ala Gly Tyr Leu Ala
 115 120 125
 Met Asp Tyr Trp Gly Gln Gly Thr Ser Val Ala Val Ser Ser Gly Gly
 130 135 140
 Gly Gly Ser Gly-Gly Gly Gly Ser Gly Gly Gly Ser Gln Ile Val
 145 150 155 160
 45 Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val
 165 170 175
 Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Thr Tyr Met His Trp Phe

- 85 -

	180	185	190
	Gln Gln Lys Pro Gly Thr Ser Pro Lys Leu Trp Ile Tyr Ser Thr Ser		
	195	200	205
5	Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly		
	210	215	220
	Thr Ser Tyr Ser Leu Thr Ile Ser Arg Met Glu Ala Glu Asp Ala Ala		
	225	230	235
	Thr Tyr Tyr Cys Gln Gln Arg Ser Thr Tyr Pro Leu Thr Phe Gly Ala		
	245	250	255
10	Gly Thr Lys Leu Glu Ile Lys Arg Glu Gln Lys Leu Ile Ser Glu Glu		
	260	265	270
	Asp Leu Asn His His His His His His		
	275	280	

15 (2) INFORMATION FOR SEQ ID NO: 45:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 72 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

TCGAGATCAA ACGGGAACAA AAATCATCT CAGAAGAAGA TCTGAATCAC CACCATCACC	60
25 ACCATTAATG AG	72

(2) INFORMATION FOR SEQ ID NO: 46:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 72 base pairs

30 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

AATTCTCATT AATGGTGGTG ATGGTGGTGA TTCAGATCTT CTTCTGAGAT GAGTTTTTGT	60
35 TCCCGTTTGA TC	72

(2) INFORMATION FOR SEQ ID NO: 47:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 864 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

- 86 -

	ATGAAATACC TATTGCCTAC GGCAGCCGCT GGATTGTTAT TACTCGCTGC CCAACCAGCC	60
	ATGGCCCAGG TCCAAGTCA GCAGCCTGGG GCTGAACTGG TGAAGCCTGG GGCTTCAGTG	120
	CAGCTGTCCT GCAAGGCTTC TGGCTACACC TTCACCGGCT ACTGGATACA CTGGGTGAAG	180
	CAGAGGCCTG GACAAGGCCT TGAGTGGATT GGAGAGGTTA ATCCTAGTAC CGGTCGTTCT	240
5	GACTACAATG AGAAGTTCAA GAACAAGGCC ACACTGACTG TAGACAAATC CTCCACCACA	300
	GCCTACATGC AACTCAGCAG CCTGACATCT GAGGACTCTG CGGTCTATTA CTGTGCAAGA	360
	GAGAGGCGCT ATGGTTACGA CGATGCTATG GACTACTGGG GCCAAGGGAC CACGGTCACC	420
	GTCTCCTCAG GTGGCGGTGG CTCGGGCGGT GGTGGGTCGG GTGGCGGCGG ATCTGACATT	480
	GAGCTCTCAC AGTCTCCATC CTCCCTGGCT GTGTCAGCAG GAGAGAAGGT CACCATGAGC	540
10	TGCAATCCA GTCAGAGTCT CCTCAACAGT AGAACCCGAA AGAACTACTT GGCTTGTTAC	600
	CAGCAGAGAC CAGGGCAGTC TCCTAACTG CTGATCTATT GGGCATCCAC TAGGACATCT	660
	GGGGTCCCTG ATCGCTTCAC AGGCAGTGA TCTGGGACAG ATTCACTCT CACCATCAGC	720
	AGTGTGCAGG CTGAAGACCT GGCAATTAT TACTGCAAGC AATCTTATAC TCTTCGGACG	780
	TTCGGTGGAG GCACCAAGCT CGAGATCAAA CGGGAACAAA AACTCATCTC AGAAGAAGAT	840
15	CTGAATCACC ACCATCACCA CCAT	864

(2) INFORMATION FOR SEQ ID NO: 48:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

25

AAGCTTGGA TTCAGTGTGA GGTGCAGCTG CAGC

34

(2) INFORMATION FOR SEQ ID NO: 49:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 45 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

35

CGCCACCTCC GGAGCCACCA CCGCCCCGTT TGATCTCGAG CTGG

45

(2) INFORMATION FOR SEQ ID NO: 50:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1998 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

45

- 87 -

ATGAAGTTGT GGCTGAACTG GATTTTCCTT GTAACACTTT TAAATGGAAT TCAGTGTGAG 60
 GTGCAGCTGC AGCAGTCTGG GGCAGAGCTT GTGAGGTCAG GGGCCTCAGT CAAGTTGTCC 120
 TGCACAGCTT CTGGCTTCAA CATTAAAGAC AACTATATGC ACTGGGTGAA GCAGAGGCCT 180
 GAACAGGGCC TGGAGTGGAT TGCATGGATT GATCCTGAGA ATGGTGATAC TGAATATGCC 240
 5 CCGAAGTTCC GGGGCAAGGC CACTTTGACT GCAGACTCAT CCTCCAACAC AGCCTACCTG 300
 CACCTCAGCA GCCTGACATC TGAGGACACT GCCGTCTATT ACTGTCATGT CCTGATCTAT 360
 GCTGGTTATT TGGCTATGGA CTACTGGGGT CAAGGAACCT CAGTCGCCGT GAGCTCGGGT 420
 GCGGCTGGCT CGGGCGGTGG TGGTTCGGGT GCGGCGGAT CTCAGATTGT GCTCACCAG 480
 TCTCCAGCAA TCATGTCTGC ATCTCCAGGG GAGAAGGTCA CCATAACCTG CAGTGCCAGC 540
 10 TCAAGTGTA CTTACATGCA CTGGTTCCAG CAGAAGCCAG GCACTTCTCC CAAACTCTGG 600
 ATTTATAGCA CATCCAACCT GGCTTCTGGA GTCCTGCTC GCTTCAGTGG CAGTGGATCT 660
 GGGACCTCTT ACTCTCTCAC AATCAGCCGA ATGGAGGCTG AAGATGCTGC CACTTATTAC 720
 TGCCAGCAAA GGAGTACTTA CCCGCTCACG TTCGGTCTG GACCAAGCT CGAGATCAAA 780
 CGGGGCGGTG GTGGCTCCGG AGGTGGCGGT AGCGGTGGCG GGGGTTCCTA GAAGCGCGAC 840
 15 AACGTGCTGT TCCAGGCAGC TACCGACGAG CAGCCGGCCG TGATCAAGAC GCTGGAGAAG 900
 CTGGTCAACA TCGAGACCGG CACCGGTGAC GCCGAGGGCA TCGCCGCTGC GGGCAACTTC 960
 CTCGAGGCGG AGCTCAAGAA CCTCGGCTTC ACGGTCACGC GAAGCAAGTC GGCCGGCCTG 1020
 GTGGTGGGCG ACAACATCGT GGGCAAGATC AAGGGCCGCG GCGGCAAGAA CCTGCTGCTG 1080
 ATGTCGCACA TGGACACCGT CTACCTCAAG GGCATTCTCG CGAAGGCCCC GTTCCGCGTC 1140
 20 GAAGGCGACA AGGCCTACGG CCCGGGCATC GCCGACGACA AGGGCGGCAA CGCGGTCATC 1200
 CTGCACACGC TCAAGCTGCT GAAGGAATAC GCGTGCAGC ACTACGGCAC CATCACCCTG 1260
 CTGTTCAACA CCGACGAGGA AAAGGGTTCC TTCGGCTCGC GCGACCTGAT CCAGGAAGAA 1320
 GCCAAGCTGG CCGACTACGT GCTCTCCTTC GAGCCACCA GCGCAGGCGA CGAAAACTC 1380
 TCGCTGGGCA CCTCGGGCAT CGCCTACGTG CAGGTCCAGA TCACCGGCAA GGCCTCGCAT 1440
 25 GCCGGCGCCG CGCCGAGCT GGGCGTGAAC GCGCTGGTCG AGGCTTCCGA CCTCGTGCTG 1500
 CGCAGCATGA ACATCGACGA CAAGGCGAAG AACCTGCGCT TCCAGTGGAC CATCGCCAAG 1560
 GCCGGCCAGG TCTCGAATC CATCCCGGCC AGCGCCACGC TGAACGCCGA CGTGCCTAC 1620
 GCGCGCAACG AGGACTTCGA CGCCGCCATG AAGACGCTGG AAGAGCGCGC GCAGCAGAAG 1680
 AAGCTGCCCC AGGCCGACGT GAAGGTGATC GTCACGCGCG GCCGCCCGGC CTTCAATGCC 1740
 30 GGCGAAGGCG GCAAGAAGCT GGTGACAAG GCGGTGGCCT ACTACAAGGA AGCCGGCGGC 1800
 ACGCTGGGCG TGAAGAGCG CACCGGCGGC GGCACCGACG CGGCCTACGC CGCGCTCTCA 1860
 GGCAAGCCAG TGATCGAGAG CCTGGGCTG CCGGGCTTCG GCTACCACAG CGACAAGGCC 1920
 GAGTACGTGG ACATCAGCGC GATTCCGCGC CGCCTGTACA TGCTGCGCG CCTGATCATG 1980
 GATCTGGGCG CCGGCAAG 1998

35

(2) INFORMATION FOR SEQ ID NO: 51:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 666 amino acids

(B) TYPE: amino acid

40

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

45

Met Lys Leu Trp Leu Asn Trp Ile Phe Leu Val Thr Leu Leu Asn Gly

1

5

10

15

Ile Gln Cys Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg

- 88 -

	20	25	30
	Ser Gly Ala Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile		
	35	40	45
5	Lys Asp Asn Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu		
	50	55	60
	Glu Trp Ile Ala Trp Ile Asp Pro Glu Asn Gly Asp Thr Glu Tyr Ala		
	65	70	75
	Pro Lys Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Ser Ser Ser Asn		
	85	90	95
10	Thr Ala Tyr Leu His Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val		
	100	105	110
	Tyr Tyr Cys His Val Leu Ile Tyr Ala Gly Tyr Leu Ala Met Asp Tyr		
	115	120	125
15	Trp Gly Gln Gly Thr Ser Val Ala Val Ser Ser Gly Gly Gly Ser		
	130	135	140
	Gly Gly Gly Gly Ser Gly Gly Gly Ser Gln Ile Val Leu Thr Gln		
	145	150	155
	Ser Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Ile Thr		
	165	170	175
20	Cys Ser Ala Ser Ser Ser Val Thr Tyr Met His Trp Phe Gln Gln Lys		
	180	185	190
	Pro Gly Thr Ser Pro Lys Leu Trp Ile Tyr Ser Thr Ser Asn Leu Ala		
	195	200	205
25	Ser Gly Val Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr		
	210	215	220
	Ser Leu Thr Ile Ser Arg Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr		
	225	230	235
	Cys Gln Gln Arg Ser Thr Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys		
	245	250	255
30	Leu Glu Ile Lys Arg Gly Gly Gly Gly Ser Gly Gly Gly Ser Gly		
	260	265	270
	Gly Gly Gly Ser Gln Lys Arg Asp Asn Val Leu Phe Gln Ala Ala Thr		
	275	280	285
35	Asp Glu Gln Pro Ala Val Ile Lys Thr Leu Glu Lys Leu Val Asn Ile		
	290	295	300
	Glu Thr Gly Thr Gly Asp Ala Glu Gly Ile Ala Ala Ala Gly Asn Phe		
	305	310	315
	Leu Glu Ala Glu Leu Lys Asn Leu Gly Phe Thr Val Thr Arg Ser Lys		
	325	330	335
40	Ser Ala Gly Leu Val Val Gly Asp Asn Ile Val Gly Lys Ile Lys Gly		
	340	345	350
	Arg Gly Gly Lys Asn Leu Leu Leu Met Ser His Met Asp Thr Val Tyr		
	355	360	365
45	Leu Lys Gly Ile Leu Ala Lys Ala Pro Phe Arg Val Glu Gly Asp Lys		
	370	375	380
	Ala Tyr Gly Pro Gly Ile Ala Asp Asp Lys Gly Gly Asn Ala Val Ile		
	385	390	395
			400

- 89 -

Leu His Thr Leu Lys Leu Leu Lys Glu Tyr Gly Val Arg Asp Tyr Gly
 405 410 415
 Thr Ile Thr Val Leu Phe Asn Thr Asp Glu Glu Lys Gly Ser Phe Gly
 420 425 430
 5 Ser Arg Asp Leu Ile Gln Glu Glu Ala Lys Leu Ala Asp Tyr Val Leu
 435 440 445
 Ser Phe Glu Pro Thr Ser Ala Gly Asp Glu Lys Leu Ser Leu Gly Thr
 450 455 460
 10 Ser Gly Ile Ala Tyr Val Gln Val Gln Ile Thr Gly Lys Ala Ser His
 465 470 475 480
 Ala Gly Ala Ala Pro Glu Leu Gly Val Asn Ala Leu Val Glu Ala Ser
 485 490 495
 Asp Leu Val Leu Arg Thr Met Asn Ile Asp Asp Lys Ala Lys Asn Leu
 500 505 510
 15 Arg Phe Gln Trp Thr Ile Ala Lys Ala Gly Gln Val Ser Asn Ile Ile
 515 520 525
 Pro Ala Ser Ala Thr Leu Asn Ala Asp Val Arg Tyr Ala Arg Asn Glu
 530 535 540
 20 Asp Phe Asp Ala Ala Met Lys Thr Leu Glu Glu Arg Ala Gln Gln Lys
 545 550 555 560
 Lys Leu Pro Glu Ala Asp Val Lys Val Ile Val Thr Arg Gly Arg Pro
 565 570 575
 Ala Phe Asn Ala Gly Glu Gly Gly Lys Lys Leu Val Asp Lys Ala Val
 580 585 590
 25 Ala Tyr Tyr Lys Glu Ala Gly Gly Thr Leu Gly Val Glu Glu Arg Thr
 595 600 605
 Gly Gly Gly Thr Asp Ala Ala Tyr Ala Ala Leu Ser Gly Lys Pro Val
 610 615 620
 30 Ile Glu Ser Leu Gly Leu Pro Gly Phe Gly Tyr His Ser Asp Lys Ala
 625 630 635 640
 Glu Tyr Val Asp Ile Ser Ala Ile Pro Arg Arg Leu Tyr Met Ala Ala
 645 650 655
 Arg Leu Ile Met Asp Leu Gly Ala Gly Lys
 660 665

- 35
- (2) INFORMATION FOR SEQ ID NO: 52:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 3217 base pairs
- (B) TYPE: nucleic acid
- 40 (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

45 GAATTCGCCG CCACTATGGA TTTCAAGTG CAGATTTTCA GCTTCCTGCT AATCAGTGCT 60
 TCAGTCATAA TGTCAGAGG ACAAACTGTT CTCTCCAGT CTCCAGCAAT CCTGTCTGCA 120
 TCTCCAGGGG AGAAGGTCAC AATGACTTGC AGGGCCAGCT CAAGTGTAACTTACATTAC 180

	TGGTACCAGC	AGAAGCCAGG	TTCCTCCCC	AAATCCTGGA	TTTATGCCAC	ATCCAACCTG	240
	GCTTCTGGAG	TCCCTGCTCG	CTTCAGTGGC	AGTGGGTCTG	GGACCTCTTA	CTCTCTCACA	300
	ATCAGCAGAG	TGGAGGCTGA	AGATGCTGCC	ACTTATTACT	GCCAACATTG	GAGTAGTAAA	360
	CCACCGACGT	TCGGTGGAGG	CACCAAGCTC	GAGATCAAAC	GGACTGTGGC	TGCACCATCT	420
5	GTCTTCATCT	TCCCGCCATC	TGATGAGCAG	TTGAAATCTG	GAAGTGCCTC	TGTTGTGTGC	480
	CTGCTGAATA	ACTTCTATCC	CAGAGAGGCC	AAAGTACAGT	GGAAGGTGGA	TAACGCCCTC	540
	CAATCGGGTA	ACTCCCAGGA	GAGTGTCACT	GAGCAGGACA	GCAAGGACAG	CACCTACAGC	600
	CTCAGCAGCA	CCCTGACGCT	GAGCAAAGCA	GACTACGAGA	AACACAAAGT	CTACGCCTGC	660
	GAAGTCACCC	ATCAGGGCCT	GAGTTCGCCC	GTCAACAAAG	GCTTCAACAG	GGGAGAGTGT	720
10	TAATAGGAGC	TCGGATCCAG	ATCTGAGCTC	CTGTAGACGT	CGACATTAAT	TCCGGTTATT	780
	TTCCACCATA	TTGCCGTCTT	TTGGCAATGT	GAGGGCCCGG	AAACCTGGCC	CTGTCTTCTT	840
	GACGAGCATT	CCTAGGGGTC	TTTCCCCTCT	CGCCAAAGGA	ATGCAAGGTC	TGTTGAATGT	900
	CGTGAAGGAA	GCAGTTCCTC	TGGAAGCTTC	TTGAAGACAA	ACAACGCTCT	TAGCGACCCCT	960
	TTGCAGGCAG	CGGAACCCCC	CACCTGGCGA	CAGGTGCCTC	TGCGGCCAAA	AGCCACGTGT	1020
15	ATAAGATACA	CCTGCAAAGG	CGGCACAACC	CCAGTGCAC	GTTGTGAGTT	GGATAGTTGT	1080
	GGAAAGAGTC	AAATGGCTCT	CCTCAAGCGT	ATTCAACAAG	GGGCTGAAGG	ATGCCCAGAA	1140
	GGTACCCCAT	TGTATGGGAT	CTGATCTGGG	GCCTCGGTGC	ACATGCTTTA	CATGTGTTTA	1200
	GTGAGGTTA	AAAAACGTCT	AGGCCCCCGG	AACCACGGGG	ACGTGGTTTT	CCTTTGAAAA	1260
	ACACGATGAT	AATACCATGG	AGTTGTGGCT	GAAGTGGATT	TTCCTTGTA	CACTTTTAAA	1320
20	TGGTATCCAG	TGTGAGGTGA	AGCTGGTGGA	GTCTGGAGGA	GGCTTGGTAC	AGCCTGGGGG	1380
	TTCTCTGAGA	CTCTCCTGTG	CAACTTCTGG	GTTACACCTC	ACTGATTACT	ACATGAACTG	1440
	GGTCCGCCAG	CCTCCAGGAA	AGGCACTTGA	GTGGTTGGGT	TTTATTGGAA	ACAAAGCTAA	1500
	TGGTTACACA	ACAGAGTACA	GTGCATCTGT	GAAGGGTCGG	TTCACCATCT	CCAGAGATAA	1560
	ATCCCAAAGC	ATCCTCTATC	TTCAAATGAA	CACCCTGAGA	GCTGAGGACA	GTGCCACTTA	1620
25	TTACTGTACA	AGAGATAGGG	GGCTACGGTT	CTACTTTGAC	TACTGGGGCC	AAGGCACCAC	1680
	TCTCACAGTG	AGCTCGGCTA	GCACCAAGGG	ACCATCGGTC	TTCCCCCTGG	CCCCCTGCTC	1740
	CAGGAGCACC	TCCGAGAGCA	CAGCCGCCCT	GGGCTGCCTG	GTCAAGGACT	ACTTCCCCGA	1800
	ACCGGTGACG	GTGTCGTGGA	ACTCAGGCGC	TCTGACCAGC	GGCGTGACA	CCTTCCCGGC	1860
	TGTCCTACAG	TCCTCAGGAC	TCTACTCCCT	CAGCAGCGTC	GTGACGGTGC	CCTCCAGCAA	1920
30	CTTCGGCACC	CAGACCTACA	CCTGCAACGT	AGATCACAAG	CCCAGCAACA	CCAAGGTGGA	1980
	CAAGACAGTT	GGCGGTGGTG	GCTCTGGTGG	TGGCGGTAGC	GGTGGCGGGG	GTTCCAGAA	2040
	GGCGGACAAC	GTGCTGTTCC	AGGCAGCTAC	CGACGAGCAG	CCGGCCGTGA	TCAAGACGCT	2100
	GGAGAAGCTG	GTCAACATCG	AGACCGGCAC	CGGTGACGCC	GAGGGCATCG	CCGCTGCGGG	2160
	CAACTTCCTC	GAGGCCGAGC	TCAAGAACCT	CGGCTTCACG	GTACGCGGAA	GCAAGTCGGC	2220
35	CGGCCTGGTG	GTGGGCGACA	ACATCGTGGG	CAAGATCAAG	GGCCGCGGCG	GCAAGAACCT	2280
	GCTGCTGATG	TCGCACATGG	ACACCGTCTA	CCTCAAGGGC	ATTCTCGCGA	AGGCCCGGTT	2340
	CCGCGTCGAA	GGCGACAAGG	CCTACGGCCC	GGGCATCGCC	GACGACAAGG	GCGGCAACGC	2400
	GGTCATCCTG	CACACGCTCA	AGCTGTGAA	GGAATACGGC	GTGCGCGACT	ACGGCACCAT	2460
	CACCGTGCTG	TTCAACACCG	ACGAGGAAAA	GGGTTCCCTC	GGCTCGCGCG	ACCTGATCCA	2520
40	GGAAGAAGCC	AAGCTGGCCG	ACTACGTGCT	CTCCTTCGAG	CCCACCAGCG	CAGGCGACGA	2580
	AAAACCTCTG	CTGGGCACCT	CGGGCATCGC	CTACGTGCAG	GTCCAGATCA	CCGGCAAGGC	2640
	CTCGCATGCC	GGCGCCGCGC	CCGAGCTGGG	CGTGAACGCG	CTGGTCGAGG	CTTCCGACCT	2700
	CGTGCTGCGC	ACGATGAACA	TCGACGACAA	GGCGAAGAAC	CTGCGCTTCC	AGTGGACCAT	2760
	CGCCAAGGCC	GGCCAGGTCT	CGAACATCAT	CCCCGCCAGC	GCCACGCTGA	ACGCCGACGT	2820
45	GCGCTACGCG	CGCAACGAGG	ACTTCGACGC	CGCCATGAAG	ACGCTGGAAG	AGCGCGCGCA	2880
	GCAGAAGAAG	CTGCCCCGAGG	CCGACGTGAA	GGTGATCGTC	ACGCGCGGCC	GCCCGGCCTT	2940
	CAATGCCGCG	GAAGGCGGCA	ACAAGCTGGT	CGACAAGGCG	GTGGCCTACT	ACAAGGAAGC	3000

- 91 -

CGGCGGCACG CTGGGCGTGG AAGAGCGCAC CGGCGGCGGC ACCGACGCGG CCTACGCCGC 3060
GCTCTCAGGC AAGCCAGTGA TCGAGAGCCT GGGCCTGCCG GGCTTCGGCT ACCACAGCGA 3120
CAAGGCCGAG TACGTGGACA TCAGCGCGAT TCCGCGCCGC CTGTACATGG CTGCGCGCCT 3180
GATCATGGAT CTGGGCGCCG GCAAGTGATA ATCTAGA 3217

5

(2) INFORMATION FOR SEQ ID NO: 53:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

15 TGGATCTGAA GCTTAACTA ACTCCATGGT GACCC

35

(2) INFORMATION FOR SEQ ID NO: 54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 61 base pairs

20 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

25

GCCACGGATC CCGCCACCTC CGGAGCCACC ACCGCCACAA TCCCTGGGCA CAATTTTCTT 60
G 61

(2) INFORMATION FOR SEQ ID NO: 55:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 94 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: other nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

CCCCAGGAAG CTTGGCGGTG GTGGCTCCGG AGGTGGCGGT AGCGGTGGCG GGGGTTCCCA 60
GAAGCGCGAC AACGTGCTGT TCCAGGCAGC TACC 94

40

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 base pairs

(B) TYPE: nucleic acid

45 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- 92 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

ATGTGCGAAT TCAGCAGCAG GTTCTTGCCG CCGCGGCCCT TGATCTTGCC C

51

5 (2) INFORMATION FOR SEQ ID NO: 57:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 732 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:16..720

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

GAATTCGCGG CCACC ATG GAT TTT CAA GTG CAG ATT TTC AGC TTC CTG CTA

51

Met Asp Phe Gln Val Gln Ile Phe Ser Phe Leu Leu

1

5

10

20 ATC AGT GCT TCA GTC ATA ATG TCC AGA GGA CAA ACT GTT CTC TCC CAG

99

Ile Ser Ala Ser Val Ile Met Ser Arg Gly Gln Thr Val Leu Ser Gln

15

20

25

TCT CCA GCA ATC CTG TCT GCA TCT CCA GGG GAG AAG GTC ACA ATG ACT

147

Ser Pro Ala Ile Leu Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr

25

30

35

40

TGC AGG GCC AGC TCA AGT GTA ACT TAC ATT CAC TGG TAC CAG CAG AAG

195

Cys Arg Ala Ser Ser Ser Val Thr Tyr Ile His Trp Tyr Gln Gln Lys

45

50

55

60

CCA GGT TCC TCC CCC AAA TCC TGG ATT TAT GCC ACA TCC AAC CTG GCT

243

30 Pro Gly Ser Ser Pro Lys Ser Trp Ile Tyr Ala Thr Ser Asn Leu Ala

65

70

75

TCT GGA GTC CCT GCT CGC TTC AGT GGC AGT GGG TCT GGG ACC TCT TAC

291

Ser Gly Val Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr

80

85

90

35 TCT CTC ACA ATC AGC AGA GTG GAG GCT GAA GAT GCT GCC ACT TAT TAC

339

Ser Leu Thr Ile Ser Arg Val Glu Ala Glu Asp Ala Ala Thr Tyr Tyr

95

100

105

TGC CAA CAT TGG AGT AGT AAA CCA CCG ACG TTC GGT GGA GGC ACC AAG

387

Cys Gln His Trp Ser Ser Lys Pro Pro Thr Phe Gly Gly Gly Thr Lys

40

110

115

120

CTG GAA ATC AAA CGG GCT GAT GCT GCA CCA ACT GTA TCC ATC TTC CCA

435

Leu Glu Ile Lys Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro

125

130

135

140

CCA TCC AGT GAG CAG TTA ACA TCT GGA GGT GCC TCA GTC GTG TGC TTC

483

45 Pro Ser Ser Glu Gln Leu Thr Ser Gly Gly Ala Ser Val Val Cys Phe

145

150

155

TTG AAC AAC TTC TAC CCC AAA GAC ATC AAT GTC AAG TGG AAG ATT GAT

531

- 93 -

Leu Asn Asn Phe Tyr Pro Lys Asp Ile Asn Val Lys Trp Lys Ile Asp
 160 165 170
 GGC AGT GAA CGA CAA AAT GGC GTC CTG AAC AGT TGG ACT GAT CAG GAC 579
 Gly Ser Glu Arg Gln Asn Gly Val Leu Asn Ser Trp Thr Asp Gln Asp
 5 175 180 185
 AGC AAA GAC AGC ACC TAC AGC ATG AGC AGC ACC CTC ACG TTG ACC AAG 627
 Ser Lys Asp Ser Thr Tyr Ser Met Ser Ser Thr Leu Thr Leu Thr Lys
 190 195 200
 GAC GAG TAT GAA CGA CAT AAC AGC TAT ACC TGT GAG GCC ACT CAC AAG 675
 10 Asp Glu Tyr Glu Arg His Asn Ser Tyr Thr Cys Glu Ala Thr His Lys
 205 210 215 220
 ACA TCA ACT TCA CCC ATT GTC AAG AGC TTC AAC AGG AAT GAG TGT 720
 Thr Ser Thr Ser Pro Ile Val Lys Ser Phe Asn Arg Asn Glu Cys
 225 230 235
 15 TAATAAGAAT TC 732

(2) INFORMATION FOR SEQ ID NO: 58:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 235 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

25 Met Asp Phe Gln Val Gln Ile Phe Ser Phe Leu Leu Ile Ser Ala Ser
 1 5 10 15
 Val Ile Met Ser Arg Gly Gln Thr Val Leu Ser Gln Ser Pro Ala Ile
 20 25 30
 Leu Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
 30 35 40 45
 Ser Ser Val Thr Tyr Ile His Trp Tyr Gln Gln Lys Pro Gly Ser Ser
 50 55 60
 Pro Lys Ser Trp Ile Tyr Ala Thr Ser Asn Leu Ala Ser Gly Val Pro
 65 70 75 80
 35 Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
 85 90 95
 Ser Arg Val Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Trp
 100 105 110
 Ser Ser Lys Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 40 115 120 125
 Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro Ser Ser Glu
 130 135 140
 Gln Leu Thr Ser Gly Gly Ala Ser Val Val Cys Phe Leu Asn Asn Phe
 145 150 155 160
 45 Tyr Pro Lys Asp Ile Asn Val Lys Trp Lys Ile Asp Gly Ser Glu Arg
 165 170 175
 Gln Asn Gly Val Leu Asn Ser Trp Thr Asp Gln Asp Ser Lys Asp Ser

10 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1974 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: other nucleic acid

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 16..1956

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

	AAGCTTGCCG	CCAC	ATG	AAG	TGG	CTG	AAC	TGG	ATT	TTC	CTT	GTA	ACA	51			
			Met	Lys	Leu	Trp	Leu	Asn	Trp	Ile	Phe	Leu	Val	Thr			
		1					5					10					
	CTT	TTA	AAT	GGT	ATC	CAG	TGT	GAG	GTG	AAG	CTG	GTG	GAG	TCT	GGA	GGA	99
25	Leu	Leu	Asn	Gly	Ile	Gln	Cys	Glu	Val	Lys	Leu	Val	Glu	Ser	Gly	Gly	
		15						20					25				
	GGC	TTG	GTA	CAG	CCT	GGG	GGT	TCT	CTG	AGA	CTC	TCC	TGT	GCA	ACT	TCT	147
	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Thr	Ser	
		30					35					40					
30	GGG	TTC	ACC	TTC	ACT	GAT	TAC	TAC	ATG	AAC	TGG	GTC	CGC	CAG	CCT	CCA	195
	Gly	Phe	Thr	Phe	Thr	Asp	Tyr	Tyr	Met	Asn	Trp	Val	Arg	Gln	Pro	Pro	
		45				50					55				60		
	GGA	AAG	GCA	CTT	GAG	TGG	TTG	GGT	TTT	ATT	GGA	AAC	AAA	GCT	AAT	GGT	243
	Gly	Lys	Ala	Leu	Glu	Trp	Leu	Gly	Phe	Ile	Gly	Asn	Lys	Ala	Asn	Gly	
35					65					70				75			
	TAC	ACA	ACA	GAG	TAC	AGT	GCA	TCT	GTG	AAG	GGT	CGG	TTC	ACC	ATC	TCC	291
	Tyr	Thr	Thr	Glu	Tyr	Ser	Ala	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	
				80					85				90				
	AGA	GAT	AAA	TCC	CAA	AGC	ATC	CTC	TAT	CTT	CAA	ATG	AAC	ACC	CTG	AGA	339
40	Arg	Asp	Lys	Ser	Gln	Ser	Ile	Leu	Tyr	Leu	Gln	Met	Asn	Thr	Leu	Arg	
		95						100					105				
	GCT	GAG	GAC	AGT	GCC	ACT	TAT	TAC	TGT	ACA	AGA	GAT	AGG	GGG	CTA	CGG	387
	Ala	Glu	Asp	Ser	Ala	Thr	Tyr	Tyr	Cys	Thr	Arg	Asp	Arg	Gly	Leu	Arg	
		110					115					120					
45	TTC	TAC	TTT	GAC	TAC	TGG	GGC	CAA	GGC	ACC	ACT	CTC	ACA	GTC	TCC	TCA	435
	Phe	Tyr	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Thr	Leu	Thr	Val	Ser	Ser	
	125					130				135					140		

- 95 -

	GCC AAA ACG ACA CCC CCA TCT GTC TAT CCA CTG GCC CCT GGA TCT GCT	483
	Ala Lys Thr Thr Pro Pro Ser Val Tyr Pro Leu Ala Pro Gly Ser Ala	
	145 150 155	
	GCC CAA ACT AAC TCC ATG GTG ACC CTG GGA TGC CTG GTC AAG GGC TAT	531
5	Ala Gln Thr Asn Ser Met Val Thr Leu Gly Cys Leu Val Lys Gly Tyr	
	160 165 170	
	TTC CCT GAG CCA GTG ACA GTG ACC TGG AAC TCT GGA TCT CTG TCC AGC	579
	Phe Pro Glu Pro Val Thr Val Thr Trp Asn Ser Gly Ser Leu Ser Ser	
	175 180 185	
10	GGT GTG CAC ACC TTC CCA GCT GTC CTG CAG TCT GAC CTC TAC ACT CTG	627
	Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Asp Leu Tyr Thr Leu	
	190 195 200	
	AGC AGC TCA GTG ACT GTC CCC TCC AGC ACC TGG CCC AGC GAG ACC GTC	675
	Ser Ser Ser Val Thr Val Pro Ser Ser Thr Trp Pro Ser Glu Thr Val	
15	205 210 215 220	
	ACC TGC AAC GTT GCC CAC CCG GCC AGC AGC ACC AAG GTG GAC AAG AAA	723
	Thr Cys Asn Val Ala His Pro Ala Ser Ser Thr Lys Val Asp Lys Lys	
	225 230 235	
	ATT GTG CCC AGG GAT TGT GGC GGT GGT GGC TCC GGA GGT GGC GGT AGC	771
20	Ile Val Pro Arg Asp Cys Gly Gly Gly Ser Gly Gly Gly Gly Ser	
	240 245 250	
	GGT GGC GGC GGT TCC CAG AAG CGC GAC AAC GTG CTG TTC CAG GCA GCT	819
	Gly Gly Gly Gly Ser Gln Lys Arg Asp Asn Val Leu Phe Gln Ala Ala	
	255 260 265	
25	ACC GAC GAG CAG CCG GCC GTG ATC AAG ACG CTG GAG AAG CTG GTC AAC	867
	Thr Asp Glu Gln Pro Ala Val Ile Lys Thr Leu Glu Lys Leu Val Asn	
	270 275 280	
	ATC GAG ACC GGC ACC GGT GAC GCC GAG GGC ATC GCC GCT GCG GGC AAC	915
	Ile Glu Thr Gly Thr Gly Asp Ala Glu Gly Ile Ala Ala Ala Gly Asn	
30	285 290 295 300	
	TTC CTC GAG GCC GAG CTC AAG AAC CTC GGC TTC ACG GTC ACG CGA AGC	963
	Phe Leu Glu Ala Glu Leu Lys Asn Leu Gly Phe Thr Val Thr Arg Ser	
	305 310 315	
	AAG TCG GCC GGC CTG GTG GTG GGC GAC AAC ATC GTG GGC AAG ATC AAG	1011
35	Lys Ser Ala Gly Leu Val Val Gly Asp Asn Ile Val Gly Lys Ile Lys	
	320 325 330	
	GGC CGC GGC GGC AAG AAC CTG CTG CTG ATG TCG CAC ATG GAC ACC GTC	1059
	Gly Arg Gly Gly Lys Asn Leu Leu Leu Met Ser His Met Asp Thr Val	
	335 340 345	
40	TAC CTC AAG GGC ATT CTC GCG AAG GCC CCG TTC CGC GTC GAA GGC GAC	1107
	Tyr Leu Lys Gly Ile Leu Ala Lys Ala Pro Phe Arg Val Glu Gly Asp	
	350 355 360	
	AAG GCC TAC GGC CCG GGC ATC GCC GAC GAC AAG GGC GGC AAC GCG GTC	1155
	Lys Ala Tyr Gly Pro Gly Ile Ala Asp Asp Lys Gly Gly Asn Ala Val	
45	365 370 375 380	
	ATC CTG CAC ACG CTC AAG CTG CTG AAG GAA TAC GGC GTG CGC GAC TAC	1203
	Ile Leu His Thr Leu Lys Leu Leu Lys Glu Tyr Gly Val Arg Asp Tyr	

- 96 -

		385		390		395	
	GGC ACC ATC ACC GTG CTG TTC AAC ACC GAC GAG GAA AAG GGT TCC TTC						1251
	Gly Thr Ile Thr Val Leu Phe Asn Thr Asp Glu Glu Lys Gly Ser Phe						
		400		405		410	
5	GGC TCG GGC GAC CTG ATC CAG GAA GAA GCC AAG CTG GCC GAC TAC GTG						1299
	Gly Ser Arg Asp Leu Ile Gln Glu Glu Ala Lys Leu Ala Asp Tyr Val						
		415		420		425	
	CTC TCC TTC GAG CCC ACC AGC GCA GGC GAC GAA AAA CTC TCG CTG GGC						1347
	Leu Ser Phe Glu Pro Thr Ser Ala Gly Asp Glu Lys Leu Ser Leu Gly						
10		430		435		440	
	ACC TCG GGC ATC GCC TAC GTG CAG GTC AAC ATC ACC GGC AAG GCC TCG						1395
	Thr Ser Gly Ile Ala Tyr Val Gln Val Asn Ile Thr Gly Lys Ala Ser						
		445		450		455	
	CAT GCC GGC GCC GCG CCC GAG CTG GGC GTG AAC GCG CTG GTC GAG GCT						1443
15	His Ala Gly Ala Ala Pro Glu Leu Gly Val Asn Ala Leu Val Glu Ala						
		465		470		475	
	TCC GAC CTC GTG CTG CGC ACG ATG AAC ATC GAC GAC AAG GCG AAG AAC						1491
	Ser Asp Leu Val Leu Arg Thr Met Asn Ile Asp Asp Lys Ala Lys Asn						
		480		485		490	
20	CTG CGC TTC AAC TGG ACC ATC GCC AAG GCC GGC AAC GTC TCG AAC ATC						1539
	Leu Arg Phe Asn Trp Thr Ile Ala Lys Ala Gly Asn Val Ser Asn Ile						
		495		500		505	
	ATC CCC GCC AGC GCC ACG CTG AAC GCC GAC GTG CGC TAC GCG CGC AAC						1587
	Ile Pro Ala Ser Ala Thr Leu Asn Ala Asp Val Arg Tyr Ala Arg Asn						
25		510		515		520	
	GAG GAC TTC GAC GCC GCC ATG AAG ACG CTG GAA GAG CGC GCG CAG CAG						1635
	Glu Asp Phe Asp Ala Ala Met Lys Thr Leu Glu Glu Arg Ala Gln Gln						
		525		530		535	
	AAG AAG CTG CCC GAG GCC GAC GTG AAG GTG ATC GTC ACG CGC GGC CGC						1683
30	Lys Lys Leu Pro Glu Ala Asp Val Lys Val Ile Val Thr Arg Gly Arg						
		545		550		555	
	CCG GCC TTC AAT GCC GGC GAA GGC GGC AAG AAG CTG GTC GAC AAG GCG						1731
	Pro Ala Phe Asn Ala Gly Glu Gly Gly Lys Lys Leu Val Asp Lys Ala						
		560		565		570	
35	GTG GCC TAC TAC AAG GAA GCC GGC GGC ACG CTG GGC GTG GAA GAG CGC						1779
	Val Ala Tyr Tyr Lys Glu Ala Gly Gly Thr Leu Gly Val Glu Glu Arg						
		575		580		585	
	ACC GGC GGC GGC ACC GAC GCG GCC TAC GCC GCG CTC TCA GGC AAG CCA						1827
	Thr Gly Gly Gly Thr Asp Ala Ala Tyr Ala Ala Leu Ser Gly Lys Pro						
40		590		595		600	
	GTG ATC GAG AGC CTG GGC CTG CCG GGC TTC GGC TAC CAC AGC GAC AAG						1875
	Val Ile Glu Ser Leu Gly Leu Pro Gly Phe Gly Tyr His Ser Asp Lys						
		605		610		615	
	GCC GAG TAC GTG GAC ATC AGC GCG ATT CCG CGC CGC CTG TAC ATG GCT						1923
45	Ala Glu Tyr Val Asp Ile Ser Ala Ile Pro Arg Arg Leu Tyr Met Ala						
		625		630		635	
	GCG CGC CTG ATC ATG GAT CTG GGC GCC GGC AAG TGATAAGAAT TCCTCGAG						1974

- 97 -

Ala Arg Leu Ile Met Asp Leu Gly Ala Gly Lys
640 645

(2) INFORMATION FOR SEQ ID NO: 60:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 647 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
 10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

Met Lys Leu Trp Leu Asn Trp Ile Phe Leu Val Thr Leu Leu Asn Gly
 1 5 10 15
 Ile Gln Cys Glu Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Gln
 15 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Thr Ser Gly Phe Thr Phe
 35 40 45
 Thr Asp Tyr Tyr Met Asn Trp Val Arg Gln Pro Pro Gly Lys Ala Leu
 50 55 60
 20 Glu Trp Leu Gly Phe Ile Gly Asn Lys Ala Asn Gly Tyr Thr Thr Glu
 65 70 75 80
 Tyr Ser Ala Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Lys Ser
 85 90 95
 Gln Ser Ile Leu Tyr Leu Gln Met Asn Thr Leu Arg Ala Glu Asp Ser
 25 100 105 110
 Ala Thr Tyr Tyr Cys Thr Arg Asp Arg Gly Leu Arg Phe Tyr Phe Asp
 115 120 125
 Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Ala Lys Thr Thr
 130 135 140
 30 Pro Pro Ser Val Tyr Pro Leu Ala Pro Gly Ser Ala Ala Gln Thr Asn
 145 150 155 160
 Ser Met Val Thr Leu Gly Cys Leu Val Lys Gly Tyr Phe Pro Glu Pro
 165 170 175
 Val Thr Val Thr Trp Asn Ser Gly Ser Leu Ser Ser Gly Val His Thr
 35 180 185 190
 Phe Pro Ala Val Leu Gln Ser Asp Leu Tyr Thr Leu Ser Ser Ser Val
 195 200 205
 Thr Val Pro Ser Ser Thr Trp Pro Ser Glu Thr Val Thr Cys Asn Val
 210 215 220
 40 Ala His Pro Ala Ser Ser Thr Lys Val Asp Lys Lys Ile Val Pro Arg
 225 230 235 240
 Asp Cys Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 245 250 255
 Ser Gln Lys Arg Asp Asn Val Leu Phe Gln Ala Ala Thr Asp Glu Gln
 45 260 265 270
 Pro Ala Val Ile Lys Thr Leu Glu Lys Leu Val Asn Ile Glu Thr Gly
 275 280 285

- 98 -

Thr Gly Asp Ala Glu Gly Ile Ala Ala Ala Gly Asn Phe Leu Glu Ala
 290 295 300
 Glu Leu Lys Asn Leu Gly Phe Thr Val Thr Arg Ser Lys Ser Ala Gly
 305 310 315 320
 5 Leu Val Val Gly Asp Asn Ile Val Gly Lys Ile Lys Gly Arg Gly Gly
 325 330 335
 Lys Asn Leu Leu Leu Met Ser His Met Asp Thr Val Tyr Leu Lys Gly
 340 345 350
 Ile Leu Ala Lys Ala Pro Phe Arg Val Glu Gly Asp Lys Ala Tyr Gly
 10 355 360 365
 Pro Gly Ile Ala Asp Asp Lys Gly Gly Asn Ala Val Ile Leu His Thr
 370 375 380
 Leu Lys Leu Leu Lys Glu Tyr Gly Val Arg Asp Tyr Gly Thr Ile Thr
 385 390 395 400
 15 Val Leu Phe Asn Thr Asp Glu Glu Lys Gly Ser Phe Gly Ser Arg Asp
 405 410 415
 Leu Ile Gln Glu Glu Ala Lys Leu Ala Asp Tyr Val Leu Ser Phe Glu
 420 425 430
 Pro Thr Ser Ala Gly Asp Glu Lys Leu Ser Leu Gly Thr Ser Gly Ile
 20 435 440 445
 Ala Tyr Val Gln Val Asn Ile Thr Gly Lys Ala Ser His Ala Gly Ala
 450 455 460
 Ala Pro Glu Leu Gly Val Asn Ala Leu Val Glu Ala Ser Asp Leu Val
 465 470 475 480
 25 Leu Arg Thr Met Asn Ile Asp Asp Lys Ala Lys Asn Leu Arg Phe Asn
 485 490 495
 Trp Thr Ile Ala Lys Ala Gly Asn Val Ser Asn Ile Ile Pro Ala Ser
 500 505 510
 Ala Thr Leu Asn Ala Asp Val Arg Tyr Ala Arg Asn Glu Asp Phe Asp
 30 515 520 525
 Ala Ala Met Lys Thr Leu Glu Glu Arg Ala Gln Gln Lys Lys Leu Pro
 530 535 540
 Glu Ala Asp Val Lys Val Ile Val Thr Arg Gly Arg Pro Ala Phe Asn
 545 550 555 560
 35 Ala Gly Glu Gly Gly Lys Lys Leu Val Asp Lys Ala Val Ala Tyr Tyr
 565 570 575
 Lys Glu Ala Gly Gly Thr Leu Gly Val Glu Glu Arg Thr Gly Gly Gly
 580 585 590
 Thr Asp Ala Ala Tyr Ala Ala Leu Ser Gly Lys Pro Val Ile Glu Ser
 40 595 600 605
 Leu Gly Leu Pro Gly Phe Gly Tyr His Ser Asp Lys Ala Glu Tyr Val
 610 615 620
 Asp Ile Ser Ala Ile Pro Arg Arg Leu Tyr Met Ala Ala Arg Leu Ile
 625 630 635 640
 45 Met Asp Leu Gly Ala Gly Lys
 645

CLAIMS

- 1 A gene construct encoding a cell targeting moiety and a heterologous prodrug
activating enzyme for use as a medicament in a mammalian host wherein the gene construct is
capable of expressing the cell targeting moiety and heterologous prodrug activating enzyme as
5 a conjugate within a cell in the mammalian host and wherein the conjugate is directed to leave
the cell thereafter for selective localisation at a cell surface antigen recognised by the cell
targeting moiety.
- 2 A gene construct for use as a medicament according to claim 1 wherein the cell
targeting moiety is an antibody.
- 10 3 A gene construct for use as a medicament according to claim 2 wherein the antibody is
an anti-CEA antibody selected from antibody A5B7 or 806.077 antibody.
- 4 A gene construct for use as a medicament according to any preceding claim wherein
the heterologous prodrug activating enzyme is a carboxypeptidase.
- 5 A gene construct for use as a medicament according to claim 4 wherein the
15 carboxypeptidase is CPG2.
- 6 A gene construct for use as a medicament according to claim 5 wherein the CPG2 has
mutated polypeptide glycosylation sites so as to prevent or reduce glycosylation on expression
in mammalian cells.
- 7 A gene construct for use as a medicament according to any one of claims 5-6 in which
20 the antibody-enzyme CPG2 conjugate is a fusion protein in which the enzyme is fused to the
C terminus of the antibody through the heavy or light chain thereof whereby dimerisation of
the encoded conjugate when expressed can take place through a dimerisation domain on
CPG2.
- 8 A gene construct for use as a medicament according to claim 7 wherein the fusion
25 protein is formed through linking a C-terminus of an antibody Fab heavy chain to an N-
terminus of a CPG2 molecule to form a Fab-CPG2 whereby two Fab-CPG2 molecules when
expressed dimerise through CPG2 to form a (Fab-CPG2)₂ conjugate.
- 9 A gene construct for use as a medicament according to claim 4 wherein the
carboxypeptidase is selected from [D253K]HCPB, [G251T,D253K]HCPB or
30 [A248S,G251T,D253K]HCPB.

- 10 A gene construct for use as a medicament according to any preceding claim comprising a transcriptional regulatory sequence which comprises a promoter and a control element which comprises a genetic switch to control expression of the gene construct.
- 11 A gene construct for use as a medicament according to claim 10 in which the
5 transcriptional regulatory sequence comprises a genetic switch control element regulated by presence of tetracycline or ecdysone.
- 12 A gene construct for use as a medicament according to claim 10 or 11 wherein the promoter is dependent on cell type and is selected from the following promoters: carcinoembryonic antigen (CEA); alpha-foetoprotein (AFP); tyrosine hydroxylase; choline
10 acetyl transferase; neurone specific enolase; insulin; glial fibro acidic protein; HER-2/neu; c-erbB2; and N-myc.
- 13 A gene construct for use as a medicament according to any preceding claim which is packaged within an adenovirus for delivery to the mammalian host.
- 14 Use of a gene construct as defined in any one of claims 1-12 for manufacture of a
15 medicament for cancer therapy in a mammalian host.
- 15 A matched two component system designed for use in a mammalian host in which the components comprise:
- (i) a first component that comprises a gene construct as defined in any one of claims 1-13 and;
 - 20 (ii) a second component that comprises a prodrug which can be converted into a cytotoxic drug by the heterologous enzyme encoded by the first component.
- 16 A matched two component system according to claim 15 in which:
the first component comprises a gene encoding the heterologous enzyme CPG2; and
the second component prodrug is selected from N-(4-[N,N-bis(2-iodoethyl)amino]-
25 phenoxy carbonyl)-L-glutamic acid, N-(4-[N,N-bis(2-chloroethyl)amino]-
phenoxy carbonyl)-L-glutamic-gamma-(3,5-dicarboxy)anilide or N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy carbonyl)-L-glutamic acid or a pharmaceutically acceptable salt thereof.
- 17 A method for the delivery of a cytotoxic drug to a site which comprises administering
30 to a host a first component that comprises a gene construct as defined in any one of claims 1-13; followed by administration to the host of a second component that comprises a prodrug

which can be converted into a cytotoxic drug by the heterologous enzyme encoded by the first component.

18 A method according to claim 17 in which the first component comprises a gene encoding the heterologous enzyme CPG2; and the second component prodrug is selected from
5 N-(4-[N,N-bis(2-iodoethyl)amino]phenoxy-carbonyl)-L-glutamic acid, N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic-gamma-(3,5-dicarboxy)anilide or N-(4-[N,N-bis(2-chloroethyl)amino]-phenoxy-carbonyl)-L-glutamic acid or a pharmaceutically acceptable salt thereof.



Figure 1**Figure 2**

Figure 3



3/3

Figure 4





Figure 5





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/62, A61K 48/00 // C07K 16/28, C12N 9/48	A3	(11) International Publication Number: WO 98/51787 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/GB98/01294 (22) International Filing Date: 5 May 1998 (05.05.98) (30) Priority Data: 9709421.3 10 May 1997 (10.05.97) GB (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): EMERY, Stephen, Charles [GB/GB]; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). BLAKEY, David, Charles [GB/GB]; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (74) Agent: GILES, Allen, Frank; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 1 April 1999 (01.04.99)	
(54) Title: GENE CONSTRUCT ENCODING A HETEROLOGOUS PRODRUG-ACTIVATING ENZYME AND A CELL TARGETING MOIETY (57) Abstract <p>The invention provides a gene construct encoding a cell targeting moiety and a heterologous prodrug activating enzyme for use as a medicament in a mammalian host wherein the gene construct is capable of expressing the cell targeting moiety and enzyme as a conjugate within a target cell in the mammalian host and wherein the conjugate is directed to leave the cell thereafter for selective localisation at a cell surface antigen recognised by the cell targeting moiety.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 98/01294

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/62 A61K48/00 //C07K16/28,C12N9/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 15341 A (CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED) 8 June 1995 see example 4 see claims	1-5,7,8, 14,15,17
Y	---	6,9,16, 18
Y	WO 97 07769 A (ZENECA LTD.) 6 March 1997 see examples see claims	6,9
Y	WO 94 02450 A (ZENECA LTD. ET AL.) 3 February 1994 see claims	16,18

	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"S" document member of the same patent family

Date of the actual completion of the international search

8 February 1999

Date of mailing of the international search report

18/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Nooij, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/01294

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	R. BEGENT ET AL.: "Single-chain Fv antibodies for targeting cancer therapy." BIOCHEMICAL SOCIETY TRANSACTIONS, vol. 25, no. 2, May 1997, pages 715-717, XP002092530 see the whole document ---	1-5,7,8, 14,15,17
X	EP 0 501 215 A (BEHRINGWERKE) 2 September 1992 see examples see claims ---	1,2,14, 15,17
X	K. BOSSLET ET AL.: "Molecular and functional characterisation of a fusion protein suited for tumour specific prodrug activation." BRITISH JOURNAL OF CANCER, vol. 65, no. 2, February 1992, pages 234-238, XP002092531 London, GB see the whole document ---	1,2,14, 15,17
X	K. BOSSLET ET AL.: "Tumor-selective prodrug activation by fusion protein-mediated catalysis." CANCER RESEARCH, vol. 54, no. 8, 15 April 1994, pages 2151-2159, XP002092532 Baltimore, MD, USA see the whole document ---	1,2,14, 15,17
A	R. SHERWOOD: "Advanced drug delivery reviews: enzyme prodrug therapy." ADVANCED DRUG DELIVERY REVIEWS, vol. 22, 1996, pages 269-288, XP000749297 Amsterdam, The Netherlands see page 270, right-hand column, line 17 - page 272, right-hand column, line 33 see page 280, right-hand column, line 24 - line 32 ---	1-18
A	H. SVENSSON ET AL.: "Synthesis and characterization of monoclonal antibody-beta-lactamase conjugates." BIOCONJUGATE CHEMISTRY, vol. 5, no. 3, May 1994, pages 262-267, XP000446101 Washington, DC, USA see abstract -----	7,8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 98/01294

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 17 and 18 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/01294

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9515341 A	08-06-1995	AU 695275 B	13-08-1998
		AU 1194795 A	19-06-1995
		CA 2177584 A	08-06-1995
		EP 0733072 A	25-09-1996
		JP 9505737 T	10-06-1997
		NZ 277057 A	27-07-1997
WO 9707769 A	06-03-1997	AU 4269796 A	19-07-1996
		AU 6747696 A	19-03-1997
		BR 9510490 A	13-01-1998
		CA 2205091 A	04-07-1996
		CA 2227040 A	06-03-1997
		CN 1171054 A	21-01-1998
		CN 1193278 A	16-09-1998
		CZ 9800430 A	13-05-1998
		CZ 9701952 A	17-12-1997
		EP 0806964 A	19-11-1997
		EP 0844885 A	03-06-1998
		FI 972683 A	19-06-1997
		WO 9620011 A	04-07-1996
		GB 2309033 A	16-07-1997
		GB 2317387 A	25-03-1998
		HU 77450 A	28-04-1998
		JP 10511547 T	10-11-1998
		NO 972882 A	19-08-1997
		NO 980630 A	07-04-1998
		PL 320964 A	24-11-1997
		PL 325017 A	06-07-1998
		SK 80997 A	04-02-1998
WO 9402450 A	03-02-1994	AT 172450 T	15-11-1998
		AU 681349 B	28-08-1997
		AU 4715693 A	14-02-1994
		CA 2101104 A	24-01-1994
		CZ 9500151 A	15-11-1995
		DE 69321729 D	26-11-1998
		EP 0651740 A	10-05-1995
		FI 950230 A	19-01-1995
		HU 69288 A	28-09-1995
		IL 106459 A	08-02-1998
		JP 7509461 T	19-10-1995
		MX 9304449 A	31-05-1994
		NO 950210 A	23-01-1995
		NZ 254864 A	26-11-1996
		PL 307226 A	15-05-1995
		SK 6995 A	11-07-1995
		US 5405990 A	11-04-1995
		US 5587161 A	24-12-1996
		US 5660829 A	26-08-1997
		US 5714148 A	03-02-1998
		ZA 9305290 A	26-04-1994
		ZW 9293 A	30-03-1994
EP 501215 A	02-09-1992	DE 4106389 A	03-09-1992
		AU 660445 B	29-06-1995
		AU 1125192 A	28-01-1993
		CA 2062047 A	29-08-1992
		JP 7179500 A	18-07-1995